

# Site report 2018 Wad Sharifey SUDAN



## Population


**21,199** refugees

**14%**

Children under 5

**20%**

Women of reproductive age



### Origin of refugees

ERITREA

ETHIOPIA

SOMALIA

NONE

Site opened  
Jan 1983

## Partners



## Key Indicators

### CRUDE MORTALITY RATE

**0.3** Deaths/1,000 refugees/month

### U5 MORTALITY RATE

**0.6** Deaths/1,000 refugees/month

### MATERNAL MORTALITY

**0** Deaths

### SKILLED BIRTH ATTENDANCE

**101%** of all births

### GAM

**9%** Global Acute Malnutrition

### SAM

**2%** Severe Acute Malnutrition


### WATER SUPPLY

**25** Liters/ person/ day



### LATRINE COVERAGE

**17** Refugees/communal toilets or latrines

## COMMUNICABLE DISEASES




## INJURIES



## NON-COMMUNICABLE DISEASES



## MENTAL HEALTH



## Public Health

HEALTH STAFFING			MORBIDITY		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Number of medical doctors		1 : < 50,000	⊖	4	
Number of qualified nurses		1 : < 10,000	⊖	24	
Number of community health workers		1 : < 1,000	⊖	145	
ACCESS AND UTILIZATION			VACCINATION		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Consultations per trained clinician per day	40	< 50	✓	Incidence of pneumonia among over 5	42
Health utilisation rate (new visits / refugee / year)	2.1	1 - 4	✓	Tuberculosis success rate	< 90%
Proportion of host population consultations	35%			Were any MDR/X-TB cases diagnosed among PoCs?	No
MORTALITY			IN-PATIENT DEPARTMENT (IPD)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.3	< 0.75	✓	Average length of stay (days)	4.3
Under-five Mortality Rate (U5MR) (/1000/month)	0.6	< 1.5	✓	Case fatality rate	0.0
Infant Mortality Rate (IMR) (/1000 livebirths)	33.8	< 30	⚠	Hospitalisation rate	32.9
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	11	< 20	✓		50 - 150
OUTBREAK ALERT AND RESPONSE					
	INDICATOR	STANDARD			
Proportion of outbreaks investigated within 48 hours		100%			
Number of outbreaks reported	0				

## Reproductive Health

ANTENATAL CARE			FAMILY PLANNING		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Antenatal care coverage	98%	> 90%	✓	18%	≥ 30%
Coverage of antenatal tetanus vaccination	98%	> 95%	✓	Source of contraceptive prevalence data	
DELIVERY CARE			SEXUAL AND GENDER-BASED VIOLENCE (SGBV)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of births attended by skilled personnel	101%	≥ 90%	✓	Total number of reported rapes	0
Proportion of births conducted by caesarean section	0%	5 - 15%	✗	Proportion of eligible rape survivors provided with PEP within 72 hours	100%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	0%	< 15%	✓	Proportion of eligible rape survivors provided with ECP within 120 hours	100%
Proportion of teenage pregnancies	0%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	
Still birth rate ( / 1000 total births / month)	0			Number of genital ulcer diseases	0
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	0
Proportion of maternal deaths investigated within 48 hours		100%			

## HIV/AIDS

### PREVENTION & TREATMENT

	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%
Condom distribution rate	0.06	> 0.5
Proportion of TB patients tested for HIV		100%

### PREVENTION & TREATMENT

	INDICATOR	STANDARD	
Number of PoCs receiving ART	0		
Proportion of HIV positive mothers who swallowed ARV during delivery		100%	✗
PMTCT coverage	82%	100%	✗

## Nutrition

### NUTRITION GENERAL

	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	9%	< 20%
Prevalence of stunting (6-59 months)	30%	< 20%
Prevalence of severe acute malnutrition (6 - 59 months)	2%	< 3%
Prevalence of anaemia in children (6-59 months)	38%	< 20%
Prevalence of anaemia in women of reproductive age (15-49 yrs)	27%	< 20%

### INFANT AND YOUNG CHILD FEEDING (IYCF)

	INDICATOR	STANDARD	
Proportion of children 0-23 months who were put to the breast within one hour of birth	93%	> 80%	✗
Proportion of infants aged between 0-5 months who received only breast milk during the previous day	91%	> 70%	✗
IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	

### MICRONUTRIENT DEFICIENCY

	INDICATOR	STANDARD
Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)		> 70%

### CMAM: SAM TREATMENT

	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	164	
Recovery rate	96%	> 75%
Default rate	0%	< 0%
Death rate	1%	< 10%
Referral rate	3%	

### MAM TREATMENT

	INDICATOR	STANDARD	
Number of new moderate acute admissions (6-59 months)	484		
Recovery rate	100%	> 75%	✓
Default rate	0%	< 15%	✓
Death rate	0%	< 3%	✓
Referral rate	0%		

## Water, Sanitation and Hygiene (WASH)

### WATER

	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	25	< 20	✓
Proportion of households collecting >= 15 litres per person per day		> 95%	
Number of persons of concern per water tap		< 80	
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual ( >= 0.1 mg/litre) and NTU <5		> 95%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%	
Proportion of households collecting drinking water from protected water sources only		> 95%	
Proportion of households with sufficient daily water storage capacity		> 80%	

### WATER STAFFING

	INDICATOR	STANDARD	
Number of hygiene promoters		1 : < 20,000	⊖

### SANITATION

	INDICATOR	STANDARD	
Refugees per latrine/toilet	17	< 20	✓
Proportion of households with drop-hole latrine or drop-hole toilet			
Proportion of households report defecating in a toilet		> 85%	
Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%	