
# Site report 2018 Kilo 26 SUDAN

## **Population**

**12,719** refugees







#### **Origin of refugees**

**ERITREA ETHIOPIA** 

**SOMALIA** 

NONE

Site opened Jan 1979

### **Partners**



### **Key Indicators**

**CRUDE MORTALITY RATE** 

Deaths/1,000 refugees/

GAM

9% Global Acute **U5 MORTALITY RATE** 

Deaths/1,000 refugees/ month

SAM

2%

Severe Acute

MATERNAL MORTALITY

Deaths

WATER SUPPLY

39

Liters/ person/ day


SKILLED BIRTH **ATTENDANCE** 


70% of all births

LATRINE COVERAGE


Refugees/communal toilets or latrines

#### **COMMUNICABLE DISEASES**





### NON-COMMUNICABLE DISEASES



**MENTAL HEALTH** 

## **Public Health**

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARD		
Number of medical doctors		1:<50,000	•	Incidence of malaria among children under	11			
Number of qualified nurses		1:<10,000	•	Incidence of watery diarrhoea among	0			
Number of community health workers		1 : < 1,000	•	children under 5				
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	62			
Consultations per trained clinician per day	29	< 50	<b>②</b>	Incidence of pneumonia among over 5	21			
Health utilisation rate (new visits / refugee / year)	1.2	1 - 4	•	Tuberculosis success rate		< 90%		
Proportion of host population consultations	42%			Were any MDR/X-TB cases diagnosed among PoCs?	No			
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATO	STAND	ARD	
Crude Mortality Rate (CMR) (/1000/month)	0.2	< 0.75	<b>②</b>	Full vaccination coverage	59%	< 95	%	8
Under-five Mortality Rate (U5MR) (/1000/month)	0.1	< 1.5	<b>②</b>	Measles vaccination coverage	59%	< 95	%	8
Infant Mortality Rate (IMR) (/1000 livebirths)	0.0	< 30	<b>②</b>	Source of vaccination data				
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	<b>②</b>	IN-PATIENT DEPARTMENT (IPD)	INDICATO	STAND	ARD	
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	6.7			
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0			
Number of outbreaks reported	0			Hospitalisation rate	97.6	50 - 1	50	<b>②</b>

# **Reproductive Health**

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	100%	> 90%	<b>Ø</b>	Contraceptive prevalence rate	8%	≥ 30%	
Coverage of antenatal tetanus vaccination	100%	> 95%	<b>Ø</b>	Source of contraceptive prevalence data			
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	70%	≥ 90%	8	Total number of reported rapes	0		
Proportion of births conducted by caesarean section	0%	5 - 15%	8	Proportion of eligible rape survivors provided with PEP within 72 hours		100%	
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	0%	< 15%	<b>②</b>	Proportion of eligible rape survivors provided with ECP within 120 hours		100%	
Proportion of teenage pregnancies	0%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate ( / 1000 total births / month)	1			Number of genital ulcer diseases	0		
Number of maternal deaths	1			Number of Pelvic Inflammatory diseases	14		
Proportion of maternal deaths investigated within 48 hours	0%	100%	8				

## HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD		PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%		Number of PoCs receving ART	0	
Condom distribution rate	0.00	> 0.5	8	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%		PMTCT coverage	4%	100%

# Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD		INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD	
Prevalence of global acute malnutrition (6-59 months)	9%	< 20%	<b>②</b>	Proportion of children 0-23 months who were put to the breast within one hour of birth	93%	> 80%	8
Prevalence of stunting (6-59 months)	30%	< 20%	<b>②</b>	Proportion of infants aged between 0-5 months who received only breast milk during the previous day	91%	> 70%	8
Prevalence of severe acute malnutrition (6 - 59 months)	2%	< 3%	<b>②</b>	IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	
Prevalence of anaemia in children (6-59 months)	38%	< 20%	8	MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	27%	< 20%	8	Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)		> 70%	
CMAM: SAM TREATMENT	INDICATOR	STANDARD		MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)	72			Number of new moderate acute admissions (6-59 months)	193		
Recovery rate	96%	> 75%	<b>②</b>	Recovery rate	94%	> 75%	<b>②</b>
Default rate	0%	< 0%	8	Default rate	0%	< 15%	<b>②</b>
Death rate	0%	< 10%	<b>Ø</b>	Death rate	0%	< 3%	<b>②</b>
Referral rate	0%			Referral rate	2%		

# Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	39	< 20	<b>②</b>	Number of hygiene promoters		1:<20,000	•
Proportion of households collecting >= 15		> 95%		SANITATION	INDICATOR	STANDARD	
litres per person per day				Refugees per latrine/toilet	12	< 20	<b>⊘</b>
Number of persons of concern per water tap		< 80		Proportion of households with drop-hole latrine or drop-hole toilet			
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual ( >= 0.1 mg/litre) and NTU <5		> 95%		Proportion of households report defecating in a toilet		> 85%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%		Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%	
Proportion of households collecting drinking water from protected water sources only		> 95%					
Proportion of households with sufficient daily water storage capacity		> 80%					