
Site report 2018 Um Gargour SUDAN

Population

13,381 refugees







Origin of refugees

ERITREA

ETHIOPIA

NONE

Site opened Jan 1976

Partners



Key Indicators

CRUDE MORTALITY RATE

Deaths/1,000 refugees/

GAM

Global Acute

U5 MORTALITY RATE

Deaths/1,000 refugees/

SAM

2%

Severe Acute Malnutrition

MATERNAL MORTALITY

Deaths

WATER SUPPLY

35

Liters/ person/ day


SKILLED BIRTH **ATTENDANCE**


58% of all births

LATRINE COVERAGE


Refugees/communal toilets or latrines

COMMUNICABLE DISEASES





NON-COMMUNICABLE DISEASES



Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STAN	NDARD	
Number of medical doctors		1:<50,000	•	Incidence of malaria among children under 5	1			
Number of qualified nurses		1:<10,000	•	Incidence of watery diarrhoea among	33			
Number of community health workers		1:<1,000	•	children under 5				
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	22			
Consultations per trained clinician per day	23	< 50	•	Incidence of pneumonia among over 5	11			
Health utilisation rate (new visits / refugee /	1.7	1 - 4	•	Tuberculosis success rate		<	90%	
year)	20/			Were any MDR/X-TB cases diagnosed among PoCs?			No	
Proportion of host population consultations	2%							
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATO	R	STANDARD	
Crude Mortality Rate (CMR) (/1000/month)	0.2	< 0.75	②	Full vaccination coverage	0%		< 95%	8
Under-five Mortality Rate (U5MR) (/1000/month)	0.3	< 1.5	②	Measles vaccination coverage	77%		< 95%	8
Infant Mortality Rate (IMR) (/1000 livebirths)	7.2	< 30	②	Source of vaccination data				
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	⊘	IN-PATIENT DEPARTMENT (IPD)	INDICATO	R	STANDARD	
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	1.6			
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0			
Number of outbreaks reported	0			Hospitalisation rate	285.9		50 - 150	②

Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD
Antenatal care coverage	76%	> 90%	8	Contraceptive prevalence rate	2%	≥ 30%
Coverage of antenatal tetanus vaccination	77%	> 95%	<u> </u>	Source of contraceptive prevalence data		
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD
Proportion of births attended by skilled personnel	58%	≥ 90%	8	Total number of reported rapes	0	
Proportion of births conducted by caesarean section	6%	5 - 15%	②	Proportion of eligible rape survivors provided with PEP within 72 hours		100%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	0%	< 15%	②	Proportion of eligible rape survivors provided with ECP within 120 hours		100%
Proportion of teenage pregnancies	0%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD
Still birth rate (/ 1000 total births / month)	19			Number of genital ulcer diseases	0	
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	0	
Proportion of maternal deaths investigated within 48 hours		100%				

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD		PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%		Number of PoCs receving ART	0	
Condom distribution rate	0.00	> 0.5	8	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%		PMTCT coverage	52%	100%

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD		INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD	
Prevalence of global acute malnutrition (6-59 months)	11%	< 20%	A	Proportion of children 0-23 months who were put to the breast within one hour of birth	83%	> 80%	8
Prevalence of stunting (6-59 months)	45%	< 20%	Ø	Proportion of infants aged between 0-5 months who received only breast milk during the previous day	81%	> 70%	8
Prevalence of severe acute malnutrition (6 - 59 months)	2%	< 3%	②	IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	
Prevalence of anaemia in children (6-59 months)	35%	< 20%	8	MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	28%	< 20%	8	Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)		> 70%	
CMAM: SAM TREATMENT	INDICATOR	STANDARD		MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)	57			Number of new moderate acute admissions (6-59 months)	280		
Recovery rate	86%	> 75%	②	Recovery rate	93%	> 75%	②
Default rate	0%	< 0%	8	Default rate	0%	< 15%	②
Death rate	7%	< 10%	②	Death rate	0%	< 3%	②
Referral rate	0%			Referral rate	4%		

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	35	< 20	•	Number of hygiene promoters		1:<20,000	•
Proportion of households collecting >= 15 litres per person per day		> 95%		SANITATION Refugees per latrine/toilet	INDICATOR 9	STANDARD < 20	©
Number of persons of concern per water tap		< 80		Proportion of households with drop-hole latrine or drop-hole toilet			
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5		> 95%		Proportion of households report defecating in a toilet		> 85%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%		Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%	
Proportion of households collecting drinking water from protected water sources only		> 95%					
Proportion of households with sufficient daily water storage capacity		> 80%					