

Site report 2018 Ajuong Thok SOUTH SUDAN



Population

37,484 refugees

18%

Children under 5

20%

Women of reproductive age



Origin of refugees

SUDAN

Site opened

Apr 2013

Partners



Key Indicators

CRUDE MORTALITY RATE

0.1 Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.3 Deaths/1,000 refugees/month

MATERNAL MORTALITY

1 Deaths

SKILLED BIRTH ATTENDANCE

96% of all births

GAM

6% Global Acute Malnutrition

SAM

0% Severe Acute Malnutrition


WATER SUPPLY

31 Liters/ person/ day


LATRINE COVERAGE

7 Refugees/communal toilets or latrines


COMMUNICABLE DISEASES





INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING			MORBIDITY		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Number of medical doctors		1 : < 50,000	⊖	51	
Number of qualified nurses		1 : < 10,000	⊖	13	
Number of community health workers		1 : < 1,000	⊖	22	
ACCESS AND UTILIZATION			VACCINATION		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Consultations per trained clinician per day	65	< 50	⊗	Incidence of pneumonia among over 5	6
Health utilisation rate (new visits / refugee / year)	1.6	1 - 4	✓	Tuberculosis success rate	< 90%
Proportion of host population consultations	19%			Were any MDR/X-TB cases diagnosed among PoCs?	No
MORTALITY			IN-PATIENT DEPARTMENT (IPD)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.1	< 0.75	✓	Average length of stay (days)	2.5
Under-five Mortality Rate (U5MR) (/1000/month)	0.3	< 1.5	✓	Case fatality rate	0.0
Infant Mortality Rate (IMR) (/1000 livebirths)	11.2	< 30	✓	Hospitalisation rate	107.1
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	2	< 20	✓		50 - 150
OUTBREAK ALERT AND RESPONSE					
	INDICATOR	STANDARD			
Proportion of outbreaks investigated within 48 hours		100%			
Number of outbreaks reported	0				

Reproductive Health

ANTENATAL CARE			FAMILY PLANNING		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Antenatal care coverage	87%	> 90%	⚠	Contraceptive prevalence rate	56%
Coverage of antenatal tetanus vaccination	89%	> 95%	⚠	Source of contraceptive prevalence data	≥ 30%
DELIVERY CARE			SEXUAL AND GENDER-BASED VIOLENCE (SGBV)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of births attended by skilled personnel	96%	≥ 90%	✓	Total number of reported rapes	10
Proportion of births conducted by caesarean section	0%	5 - 15%	⊗	Proportion of eligible rape survivors provided with PEP within 72 hours	90%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	0%	< 15%	✓	Proportion of eligible rape survivors provided with ECP within 120 hours	70%
Proportion of teenage pregnancies	6%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	
Still birth rate (/ 1000 total births / month)	1			Number of genital ulcer diseases	141
Number of maternal deaths	1			Number of Pelvic Inflammatory diseases	348
Proportion of maternal deaths investigated within 48 hours	100%	100%	✓		

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD	PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV	106%	100%	✗ Number of PoCs receiving ART	48	
Condom distribution rate	0.13	> 0.5	✗ Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%	PMTCT coverage	100%	100% ⚠

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD	INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	6%	< 20%	✓ Proportion of children 0-23 months who were put to the breast within one hour of birth	93%	> 80% ✗
Prevalence of stunting (6-59 months)	29%	< 20%	✓ Proportion of infants aged between 0-5 months who received only breast milk during the previous day	94%	> 70% ✗
Prevalence of severe acute malnutrition (6 - 59 months)	0%	< 3%	✓ IYCF programmes targeting pregnant and lactating women established or maintained?		Yes
Prevalence of anaemia in children (6-59 months)	46%	< 20%	✗ MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD
Prevalence of anaemia in women of reproductive age (15-49 yrs)	38%	< 20%	✗ Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)		> 70%

CMAM: SAM TREATMENT	INDICATOR	STANDARD	MAM TREATMENT	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	215		Number of new moderate acute admissions (6-59 months)	1009	
Recovery rate	86%	> 75%	✓ Recovery rate	88%	> 75% ✓
Default rate	2%	< 0%	✗ Default rate	1%	< 15% ✓
Death rate	2%	< 10%	✓ Death rate	0%	< 3% ✓
Referral rate	2%		Referral rate	7%	

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD	WATER STAFFING	INDICATOR	STANDARD
Average number of litres of potable water available per person per day	31	< 20	✓ Number of hygiene promoters		1 : < 20,000 -
Proportion of households collecting >= 15 litres per person per day		> 95%	SANITATION	INDICATOR	STANDARD
Number of persons of concern per water tap		< 80	Refugees per latrine/toilet	7	< 20 ✓
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5		> 95%	Proportion of households with drop-hole latrine or drop-hole toilet		> 85%
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%	Proportion of households report defecating in a toilet		> 90%
Proportion of households collecting drinking water from protected water sources only		> 95%	Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		
Proportion of households with sufficient daily water storage capacity		> 80%			