

# Site report 2018 Doro SOUTH SUDAN



## Population


**58,808** refugees

**22%**

Children under 5

**20%**

Women of reproductive age



Origin of refugees

**SUDAN**

Site opened  
**Nov 2011**

## Partners



## Key Indicators

CRUDE MORTALITY RATE

**0.1** Deaths/1,000 refugees/month

U5 MORTALITY RATE

**0.2** Deaths/1,000 refugees/month

MATERNAL MORTALITY

**4** Deaths

SKILLED BIRTH ATTENDANCE

**100%** of all births

GAM

**NC** Global Acute Malnutrition

SAM

**NC** Severe Acute Malnutrition


WATER SUPPLY

**18** Liters/ person/ day


LATRINE COVERAGE

**16** Refugees/communal toilets or latrines


### COMMUNICABLE DISEASES




### NON-COMMUNICABLE DISEASES



### INJURIES



### MENTAL HEALTH



## Public Health

### HEALTH STAFFING

	INDICATOR	STANDARD
Number of medical doctors		1 : < 50,000
Number of qualified nurses		1 : < 10,000
Number of community health workers		1 : < 1,000

### ACCESS AND UTILIZATION

	INDICATOR	STANDARD
Consultations per trained clinician per day	55	< 50
Health utilisation rate (new visits / refugee / year)	0.6	1 - 4
Proportion of host population consultations	1%	

### MORTALITY

	INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.1	< 0.75
Under-five Mortality Rate (U5MR) (/1000/month)	0.2	< 1.5
Infant Mortality Rate (IMR) (/1000 livebirths)	9.3	< 30
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	3	< 20

### OUTBREAK ALERT AND RESPONSE

	INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours		100%
Number of outbreaks reported	0	

### MORBIDITY

	INDICATOR	STANDARD
Incidence of malaria among children under 5	5	
Incidence of watery diarrhoea among children under 5	20	
Incidence of pneumonia among children under 5	34	
Incidence of pneumonia among over 5	11	
Tuberculosis success rate		< 90%
Were any MDR/X-TB cases diagnosed among PoCs?		No

### VACCINATION

	INDICATOR	STANDARD
Full vaccination coverage	28%	< 95%
Measles vaccination coverage	30%	< 95%
Source of vaccination data		

### IN-PATIENT DEPARTMENT (IPD)

	INDICATOR	STANDARD
Average length of stay (days)	0.0	
Case fatality rate	0.0	
Hospitalisation rate	6.5	50 - 150

## Reproductive Health

### ANTENATAL CARE

	INDICATOR	STANDARD
Antenatal care coverage	99%	> 90%
Coverage of antenatal tetanus vaccination	99%	> 95%

### DELIVERY CARE

	INDICATOR	STANDARD
Proportion of births attended by skilled personnel	100%	≥ 90%
Proportion of births conducted by caesarean section	0%	5 - 15%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	0%	< 15%
Proportion of teenage pregnancies	0%	
Still birth rate ( / 1000 total births / month)	0	
Number of maternal deaths	4	
Proportion of maternal deaths investigated within 48 hours	80%	100%

### FAMILY PLANNING

	INDICATOR	STANDARD
Contraceptive prevalence rate	0%	≥ 30%
Source of contraceptive prevalence data		

### SEXUAL AND GENDER-BASED VIOLENCE (SGBV)

	INDICATOR	STANDARD
Total number of reported rapes	0	
Proportion of eligible rape survivors provided with PEP within 72 hours		100%
Proportion of eligible rape survivors provided with ECP within 120 hours		100%

### SEXUALLY TRANSMITTED INFECTIONS (STIS)

	INDICATOR	STANDARD
Number of genital ulcer diseases	177	
Number of Pelvic Inflammatory diseases	133	

## HIV/AIDS

### PREVENTION & TREATMENT

Proportion of blood units screened for HIV

INDICATOR STANDARD

100%

Condom distribution rate

0.00

> 0.5

Proportion of TB patients tested for HIV

100%

### PREVENTION & TREATMENT

Number of PoCs receiving ART

0

✘ Proportion of HIV positive mothers who swallowed ARV during delivery

100%

PMTCT coverage

24%

100%



## Nutrition

### NUTRITION GENERAL

Prevalence of global acute malnutrition (6-59 months)

Not collected

Prevalence of stunting (6-59 months)

Not collected

Prevalence of severe acute malnutrition (6 - 59 months)

Not collected

Prevalence of anaemia in children (6-59 months)

Not collected

Prevalence of anaemia in women of reproductive age (15-49 yrs)

Not collected

### INFANT AND YOUNG CHILD FEEDING (IYCF)

Proportion of children 0-23 months who were put to the breast within one hour of birth

Not collected

Proportion of infants aged between 0-5 months who received only breast milk during the previous day

Not collected

IYCF programmes targeting pregnant and lactating women established or maintained?

Yes

### MICRONUTRIENT DEFICIENCY

Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)

> 70%

### CMAM: SAM TREATMENT

Number of new severe acute admissions (6-59 months)

318

Recovery rate

70%

> 75%

Default rate

8%

< 0%

Death rate

0%

< 10%

Referral rate

22%

### MAM TREATMENT

Number of new moderate acute admissions (6-59 months)

1032

⚠ Recovery rate

75%

> 75%



✘ Default rate

9%

< 15%



✔ Death rate

0%

< 3%



Referral rate

14%

## Water, Sanitation and Hygiene (WASH)

### WATER

Average number of litres of potable water available per person per day

18

< 20



Proportion of households collecting >= 15 litres per person per day

> 95%

Number of persons of concern per water tap

< 80

Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual ( >= 0.1 mg/litre) and NTU <5

> 95%

Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml

> 95%

Proportion of households collecting drinking water from protected water sources only

> 95%

Proportion of households with sufficient daily water storage capacity

> 80%

### WATER STAFFING

Number of hygiene promoters

1 : < 20,000



### SANITATION

Refugees per latrine/toilet

16

< 20



Proportion of households with drop-hole latrine or drop-hole toilet

Proportion of households report defecating in a toilet

> 85%

Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)

> 90%