
# Site report 2018 Gendrassa SOUTH SUDAN

### **Population**

**17,427** refugees





**Origin of refugees** 

Site opened Jun 2012



#### **Partners**





### **Key Indicators**

**CRUDE MORTALITY RATE** 

Deaths/1,000 refugees/

**GAM** 

NC Global Acute U5 MORTALITY RATE

Deaths/1,000 refugees/

SAM

NC

Severe Acute Malnutrition

MATERNAL MORTALITY

Deaths

WATER SUPPLY

22

Liters/ person/ day

SKILLED BIRTH **ATTENDANCE** 


40% of all births


LATRINE COVERAGE


Refugees/communal toilets or latrines


### COMMUNICABLE DISEASES

#### **NON-COMMUNICABLE DISEASES**









## Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARD		
Number of medical doctors		1:<50,000	•	Incidence of malaria among children under 5	38			
Number of qualified nurses		1:<10,000	•	Incidence of watery diarrhoea among	43			
Number of community health workers		1:<1,000	•	children under 5				
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	124			
Consultations per trained clinician per day	49	< 50	<b>②</b>	Incidence of pneumonia among over 5	38			
Health utilisation rate (new visits / refugee /	2.6	1 - 4	•	Tuberculosis success rate		< 90%		
year) Proportion of host population consultations	1%			Were any MDR/X-TB cases diagnosed among PoCs?		No		
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATO	STAN	IDARD	
Crude Mortality Rate (CMR) (/1000/month)	0.1	< 0.75	<b>⊘</b>	Full vaccination coverage	72%	< 9	95%	8
Under-five Mortality Rate (U5MR) (/1000/month)	0.4	< 1.5	<b>②</b>	Measles vaccination coverage	77%	< 9	95%	3
Infant Mortality Rate (IMR) (/1000 livebirths)	21.0	< 30	<b>⊘</b>	Source of vaccination data				
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	4	< 20	<b>②</b>	IN-PATIENT DEPARTMENT (IPD)	INDICATO	STAN	IDARD	
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	0.0			
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0			
Number of outbreaks reported	0			Hospitalisation rate	29.4	50 -	- 150	8

# Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	77%	> 90%	8	Contraceptive prevalence rate	47%	≥ 30%	•
Coverage of antenatal tetanus vaccination	92%	> 95%	<b>A</b>	Source of contraceptive prevalence data			
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	40%	≥ 90%	8	Total number of reported rapes	4		
Proportion of births conducted by caesarean section	0%	5 - 15%	8	Proportion of eligible rape survivors provided with PEP within 72 hours	100%	100%	•
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	2%	< 15%	<b>②</b>	Proportion of eligible rape survivors provided with ECP within 120 hours	0%	100%	×
Proportion of teenage pregnancies	1%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate ( / 1000 total births / month)	2			Number of genital ulcer diseases	65		
Number of maternal deaths	1			Number of Pelvic Inflammatory diseases	167		
Proportion of maternal deaths investigated within 48 hours	100%	100%	<b>⊘</b>				

## HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD		PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%		Number of PoCs receving ART	0	
Condom distribution rate	0.01	> 0.5	8	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%		PMTCT coverage	97%	100%

## Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD		INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD	
Prevalence of global acute malnutrition (6-59 months)	Not collected			Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected		
Prevalence of stunting (6-59 months)	Not collected			Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected		
Prevalence of severe acute malnutrition (6 - 59 months)	Not collected			IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	
Prevalence of anaemia in children (6-59 months)	Not collected			MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected			Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)		> 70%	
CMAM: SAM TREATMENT	INDICATOR	STANDARD		MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)	163			Number of new moderate acute admissions (6-59 months)	355		
Recovery rate	82%	> 75%	<b>②</b>	Recovery rate	81%	> 75%	<b>②</b>
Default rate	0% < 0%		8	Default rate	0%	< 15%	<b>②</b>
Death rate	0%	< 10%	<b>②</b>	Death rate	0%	< 3%	
Referral rate	18%			Referral rate	14%		

# Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	22	< 20	•	Number of hygiene promoters		1:<20,000	•
Proportion of households collecting >= 15 litres per person per day		> 95%		SANITATION  Polygood pay latring /tailet	INDICATOR 8	STANDARD < 20	<b>•</b>
Number of persons of concern per water tap		< 80		Refugees per latrine/toilet  Proportion of households with drop-hole	0	< 20	
Proportion of tests at chlorinated water collection points indicating a measurable		> 95%		latrine or drop-hole toilet		. 050/	
chlorine residual ( >= 0.1 mg/litre) and NTU <5				Proportion of households report defecating in a toilet		> 85%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%		Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%	
Proportion of households collecting drinking water from protected water sources only		> 95%					
Proportion of households with sufficient daily water storage capacity		> 80%					