
Site report 2018 Gentil Hospital SOUTH SUDAN

Population







Site opened **Invalid date**

UNHCR The UN Refugee Agency

Partners

Key Indicators

CRUDE MORTALITY RATE

1328.0

GAM

Deaths/1,000 refugees/

U5 MORTALITY RATE

Deaths/1,000 refugees/

MATERNAL MORTALITY

Deaths

WATER SUPPLY

SKILLED BIRTH ATTENDANCE

92% of all births

LATRINE COVERAGE

Refugees/communal toilets or latrines

Global Acute

NC


SAM

Severe Acute Malnutrition


Liters/ person/ day

COMMUNICABLE DISEASES


45.6 %



harm



NON-COMMUNICABLE DISEASES



Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARD
Number of medical doctors		1:<50,000	•	Incidence of malaria among children under 5		
Number of qualified nurses		1:<10,000	•	Incidence of watery diarrhoea among		
Number of community health workers		1:<1,000	•	children under 5		
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5		
Consultations per trained clinician per day	26	< 50	•	Incidence of pneumonia among over 5	89,848	
Health utilisation rate (new visits / refugee /	15311.5	1 - 4	•	Tuberculosis success rate		< 90%
year) Proportion of host population consultations	10%			Were any MDR/X-TB cases diagnosed among PoCs?		No
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	1328.0	< 0.75	8	Full vaccination coverage		< 95%
Under-five Mortality Rate (U5MR) (/1000/month)		< 1.5		Measles vaccination coverage		< 95%
Infant Mortality Rate (IMR) (/1000 livebirths)	11.6	< 30	②	Source of vaccination data		
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	3	< 20	②	IN-PATIENT DEPARTMENT (IPD)	INDICATOR	STANDARD
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	3.3	
Proportion of outbreaks investigated within 48 hours	0%	100%	8	Case fatality rate	0.0	
Number of outbreaks reported	15			Hospitalisation rate	1219000.0	50 - 150

Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD
Antenatal care coverage	58%	> 90%	8	Contraceptive prevalence rate	88000%	≥ 30%
Coverage of antenatal tetanus vaccination	79%	> 95%	A	Source of contraceptive prevalence data		
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD
Proportion of births attended by skilled personnel	92%	≥ 90%	②	Total number of reported rapes	0	
Proportion of births conducted by caesarean section	0%	5 - 15%	8	Proportion of eligible rape survivors provided with PEP within 72 hours		100%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	6%	< 15%	②	Proportion of eligible rape survivors provided with ECP within 120 hours		100%
Proportion of teenage pregnancies	9%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD
Still birth rate (/ 1000 total births / month)	1			Number of genital ulcer diseases	24	
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	59	
Proportion of maternal deaths investigated within 48 hours		100%				

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD	PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%	Number of PoCs receving ART	0	
Condom distribution rate	54.61	> 0.5	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%	PMTCT coverage	67%	100%

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD		INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD	
Prevalence of global acute malnutrition (6-59 months)	Not collected			Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected		
Prevalence of stunting (6-59 months)	Not collected			Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected		
Prevalence of severe acute malnutrition (6 - 59 months)	Not collected			IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	
Prevalence of anaemia in children (6-59 months)	Not collected			MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected			Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)		> 70%	
CMAM: SAM TREATMENT	INDICATOR	STANDARD		MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)	135			Number of new moderate acute admissions (6-59 months)	0		
Recovery rate	12%	> 75%	8	Recovery rate	0%	> 75%	8
Default rate	35%	< 0%	A	Default rate	100%	< 15%	8
Death rate	47%	< 10%	8	Death rate	0%	< 3%	②
Referral rate	6%			Referral rate	0%		

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD	WATER STAFFING	INDICATOR	STANDARD
Average number of litres of potable water available per person per day		< 20	Number of hygiene promoters		1:<20,000
Proportion of households collecting >= 15		> 95%	SANITATION	INDICATOR	STANDARD
litres per person per day		< 80	Refugees per latrine/toilet		< 20
Number of persons of concern per water tap Proportion of tests at chlorinated water		> 95%	Proportion of households with drop-hole latrine or drop-hole toilet		
collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5		> 5570	Proportion of households report defecating in a toilet		> 85%
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%	Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%
Proportion of households collecting drinking water from protected water sources only		> 95%			
Proportion of households with sufficient daily water storage capacity		> 80%			