

Site report 2018 Gorom SOUTH SUDAN



Population


2,134 refugees

16%

Children under 5

20%

Women of reproductive age



Origin of refugees

ETHIOPIA

Site opened

Nov 2010

Partners



Key Indicators

CRUDE MORTALITY RATE

0.1

Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.2

Deaths/1,000 refugees/month

MATERNAL MORTALITY

0

Deaths

SKILLED BIRTH ATTENDANCE

92%

of all births

GAM

4%

Global Acute Malnutrition

SAM

0

Severe Acute Malnutrition


WATER SUPPLY

Liters/ person/ day


LATRINE COVERAGE

Refugees/communal toilets or latrines


COMMUNICABLE DISEASES





INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING			MORBIDITY				
	INDICATOR	STANDARD		INDICATOR	STANDARD		
Number of medical doctors		1 : < 50,000	⊖	97			
Number of qualified nurses		1 : < 10,000	⊖	27			
Number of community health workers		1 : < 1,000	⊖	33			
ACCESS AND UTILIZATION			VACCINATION				
	INDICATOR	STANDARD		INDICATOR	STANDARD		
Consultations per trained clinician per day	33	< 50	✓	Incidence of pneumonia among over 5	13		
Health utilisation rate (new visits / refugee / year)	3.9	1 - 4	✓	Tuberculosis success rate	< 90%		
Proportion of host population consultations	39%			Were any MDR/X-TB cases diagnosed among PoCs?	No		
MORTALITY			IN-PATIENT DEPARTMENT (IPD)				
	INDICATOR	STANDARD		INDICATOR	STANDARD		
Crude Mortality Rate (CMR) (/1000/month)	0.1	< 0.75	✓	Average length of stay (days)	1.5		
Under-five Mortality Rate (U5MR) (/1000/month)	0.2	< 1.5	✓	Case fatality rate	0.0		
Infant Mortality Rate (IMR) (/1000 livebirths)	0.0	< 30	✓	Hospitalisation rate	102.6	50 - 150	✓
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	✓				
OUTBREAK ALERT AND RESPONSE							
	INDICATOR	STANDARD					
Proportion of outbreaks investigated within 48 hours		100%					
Number of outbreaks reported	0						

Reproductive Health

ANTENATAL CARE			FAMILY PLANNING			
	INDICATOR	STANDARD		INDICATOR	STANDARD	
Antenatal care coverage	92%	> 90%	✓	5173%	≥ 30%	✓
Coverage of antenatal tetanus vaccination	92%	> 95%	⚠	Source of contraceptive prevalence data		
DELIVERY CARE			SEXUAL AND GENDER-BASED VIOLENCE (SGBV)			
	INDICATOR	STANDARD		INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	92%	≥ 90%	✓	Total number of reported rapes	0	
Proportion of births conducted by caesarean section	0%	5 - 15%	✗	Proportion of eligible rape survivors provided with PEP within 72 hours	100%	
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	0%	< 15%	✓	Proportion of eligible rape survivors provided with ECP within 120 hours	100%	
Proportion of teenage pregnancies	5%			SEXUALLY TRANSMITTED INFECTIONS (STIS)		
Still birth rate (/ 1000 total births / month)	0			Number of genital ulcer diseases	11	
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	6	
Proportion of maternal deaths investigated within 48 hours		100%				

HIV/AIDS

PREVENTION & TREATMENT

	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%
Condom distribution rate	0.60	> 0.5
Proportion of TB patients tested for HIV		100%

PREVENTION & TREATMENT

	INDICATOR	STANDARD	
Number of PoCs receiving ART	54		
Proportion of HIV positive mothers who swallowed ARV during delivery		100%	✓
PMTCT coverage	89%	100%	✗

Nutrition

NUTRITION GENERAL

	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	4%	< 20%
Prevalence of stunting (6-59 months)	17%	< 20%
Prevalence of severe acute malnutrition (6 - 59 months)	0%	< 3%
Prevalence of anaemia in children (6-59 months)	55%	< 20%
Prevalence of anaemia in women of reproductive age (15-49 yrs)	57%	< 20%

INFANT AND YOUNG CHILD FEEDING (IYCF)

	INDICATOR	STANDARD	
Proportion of children 0-23 months who were put to the breast within one hour of birth	117%	> 80%	✗
Proportion of infants aged between 0-5 months who received only breast milk during the previous day	93%	> 70%	✗
IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	

MICRONUTRIENT DEFICIENCY

	INDICATOR	STANDARD
Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)		> 70%

CMAM: SAM TREATMENT

	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	5	
Recovery rate	83%	> 75%
Default rate	17%	< 0%
Death rate	0%	< 10%
Referral rate	0%	

MAM TREATMENT

	INDICATOR	STANDARD	
Number of new moderate acute admissions (6-59 months)	31		
Recovery rate	81%	> 75%	✓
Default rate	3%	< 15%	✓
Death rate	0%	< 3%	✓
Referral rate	0%		

Water, Sanitation and Hygiene (WASH)

WATER

	INDICATOR	STANDARD
Average number of litres of potable water available per person per day		< 20
Proportion of households collecting >= 15 litres per person per day		> 95%
Number of persons of concern per water tap		< 80
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5		> 95%
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%
Proportion of households collecting drinking water from protected water sources only		> 95%
Proportion of households with sufficient daily water storage capacity		> 80%

WATER STAFFING

	INDICATOR	STANDARD
Number of hygiene promoters		1 : < 20,000

SANITATION

	INDICATOR	STANDARD
Refugees per latrine/toilet		< 20
Proportion of households with drop-hole latrine or drop-hole toilet		
Proportion of households report defecating in a toilet		> 85%
Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%