

Site report 2018 Kaya SOUTH SUDAN



Population


22,904 refugees

21%

Children under 5

20%

Women of reproductive age



Origin of refugees

SUDAN

Site opened
Jun 2012

Partners



ACTED
Agency for Technical Cooperation and Development

Key Indicators

CRUDE MORTALITY RATE

0.1

Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.4

Deaths/1,000 refugees/month

MATERNAL MORTALITY

0

Deaths

SKILLED BIRTH ATTENDANCE

66%

of all births

GAM

NC

Global Acute Malnutrition

SAM

NC

Severe Acute Malnutrition

WATER SUPPLY

19


Liters/ person/ day

LATRINE COVERAGE


12

Refugees/communal toilets or latrines


COMMUNICABLE DISEASES




INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING			MORBIDITY				
	INDICATOR	STANDARD		INDICATOR	STANDARD		
Number of medical doctors		1 : < 50,000	⊖	17			
Number of qualified nurses		1 : < 10,000	⊖	50			
Number of community health workers		1 : < 1,000	⊖	188			
ACCESS AND UTILIZATION			VACCINATION				
	INDICATOR	STANDARD		INDICATOR	STANDARD		
Consultations per trained clinician per day	63	< 50	⊗	Incidence of pneumonia among over 5	62		
Health utilisation rate (new visits / refugee / year)	2.4	1 - 4	✓	Tuberculosis success rate	< 90%		
Proportion of host population consultations	0%			Were any MDR/X-TB cases diagnosed among PoCs?	No		
MORTALITY			IN-PATIENT DEPARTMENT (IPD)				
	INDICATOR	STANDARD		INDICATOR	STANDARD		
Crude Mortality Rate (CMR) (/1000/month)	0.1	< 0.75	✓	Average length of stay (days)	0.2		
Under-five Mortality Rate (U5MR) (/1000/month)	0.4	< 1.5	✓	Case fatality rate	0.0		
Infant Mortality Rate (IMR) (/1000 livebirths)	22.3	< 30	✓	Hospitalisation rate	13.8	50 - 150	⊗
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	2	< 20	✓				
OUTBREAK ALERT AND RESPONSE							
	INDICATOR	STANDARD					
Proportion of outbreaks investigated within 48 hours		100%					
Number of outbreaks reported	0						

Reproductive Health

ANTENATAL CARE			FAMILY PLANNING				
	INDICATOR	STANDARD		INDICATOR	STANDARD		
Antenatal care coverage	77%	> 90%	⊗	11%	≥ 30%	⊗	
Coverage of antenatal tetanus vaccination	81%	> 95%	⚠	Source of contraceptive prevalence data			
DELIVERY CARE			SEXUAL AND GENDER-BASED VIOLENCE (SGBV)				
	INDICATOR	STANDARD		INDICATOR	STANDARD		
Proportion of births attended by skilled personnel	66%	≥ 90%	⊗	Total number of reported rapes	4		
Proportion of births conducted by caesarean section	0%	5 - 15%	⊗	Proportion of eligible rape survivors provided with PEP within 72 hours	100%	100%	✓
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	3%	< 15%	✓	Proportion of eligible rape survivors provided with ECP within 120 hours	0%	100%	⊗
Proportion of teenage pregnancies	1%			SEXUALLY TRANSMITTED INFECTIONS (STIS)			
Still birth rate (/ 1000 total births / month)	3			Number of genital ulcer diseases	122		
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	561		
Proportion of maternal deaths investigated within 48 hours		100%					

HIV/AIDS

PREVENTION & TREATMENT

Proportion of blood units screened for HIV

INDICATOR STANDARD

100%

Condom distribution rate

0.01 > 0.5

Proportion of TB patients tested for HIV

100%

PREVENTION & TREATMENT

Number of PoCs receiving ART

INDICATOR STANDARD

1

✘ Proportion of HIV positive mothers who swallowed ARV during delivery

100%

PMTCT coverage

88% 100%



Nutrition

NUTRITION GENERAL

Prevalence of global acute malnutrition (6-59 months)

INDICATOR STANDARD

Not collected

Prevalence of stunting (6-59 months)

Not collected

Prevalence of severe acute malnutrition (6 - 59 months)

Not collected

Prevalence of anaemia in children (6-59 months)

Not collected

Prevalence of anaemia in women of reproductive age (15-49 yrs)

Not collected

INFANT AND YOUNG CHILD FEEDING (IYCF)

Proportion of children 0-23 months who were put to the breast within one hour of birth

INDICATOR STANDARD

Not collected

Proportion of infants aged between 0-5 months who received only breast milk during the previous day

Not collected

IYCF programmes targeting pregnant and lactating women established or maintained?

Yes

MICRONUTRIENT DEFICIENCY

Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)

INDICATOR STANDARD

> 70%

CMAM: SAM TREATMENT

Number of new severe acute admissions (6-59 months)

INDICATOR STANDARD

131

Recovery rate

65% > 75%

Default rate

0% < 0%

Death rate

0% < 10%

Referral rate

34%

MAM TREATMENT

Number of new moderate acute admissions (6-59 months)

INDICATOR STANDARD

511

✘ Recovery rate

75% > 75%



✘ Default rate

0% < 15%



✔ Death rate

0% < 3%



Referral rate

15%

Water, Sanitation and Hygiene (WASH)

WATER

Average number of litres of potable water available per person per day

INDICATOR STANDARD

19 < 20



Proportion of households collecting >= 15 litres per person per day

> 95%

Number of persons of concern per water tap

< 80

Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5

> 95%

Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml

> 95%

Proportion of households collecting drinking water from protected water sources only

> 95%

Proportion of households with sufficient daily water storage capacity

> 80%

WATER STAFFING

Number of hygiene promoters

INDICATOR STANDARD

1 : < 20,000



SANITATION

Refugees per latrine/toilet

INDICATOR STANDARD

12 < 20



Proportion of households with drop-hole latrine or drop-hole toilet

Proportion of households report defecating in a toilet

> 85%

Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)

> 90%