



Site report 2018 Makpandu SOUTH SUDAN



Population

4,057 refugees



Origin of refugees

DRC CAR ETHIOPIA SOMALIA

Site opened
Sep 2008

Partners



Key Indicators

CRUDE MORTALITY RATE

0.0 Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.1 Deaths/1,000 refugees/month

MATERNAL MORTALITY

0 Deaths

SKILLED BIRTH ATTENDANCE

96% of all births

GAM

5% Global Acute Malnutrition

SAM

1% Severe Acute Malnutrition


WATER SUPPLY

Liters/ person/ day


LATRINE COVERAGE

Refugees/communal toilets or latrines


COMMUNICABLE DISEASES





INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



HIV/AIDS

PREVENTION & TREATMENT

	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%
Condom distribution rate	0.23	> 0.5
Proportion of TB patients tested for HIV		100%

PREVENTION & TREATMENT

	INDICATOR	STANDARD	
Number of PoCs receiving ART	73		
Proportion of HIV positive mothers who swallowed ARV during delivery		100%	✗
PMTCT coverage	103%	100%	✓

Nutrition

NUTRITION GENERAL

	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	5%	< 20%
Prevalence of stunting (6-59 months)	21%	< 20%
Prevalence of severe acute malnutrition (6 - 59 months)	1%	< 3%
Prevalence of anaemia in children (6-59 months)	Not collected	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected	

INFANT AND YOUNG CHILD FEEDING (IYCF)

	INDICATOR	STANDARD
Proportion of children 0-23 months who were put to the breast within one hour of birth		Not collected
Proportion of infants aged between 0-5 months who received only breast milk during the previous day		Not collected
IYCF programmes targeting pregnant and lactating women established or maintained?		Yes

MICRONUTRIENT DEFICIENCY

	INDICATOR	STANDARD
Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/BBF)		> 70%

CMAM: SAM TREATMENT

	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	14	
Recovery rate	83%	> 75%
Default rate	8%	< 0%
Death rate	0%	< 10%
Referral rate	0%	

MAM TREATMENT

	INDICATOR	STANDARD	
Number of new moderate acute admissions (6-59 months)	39		
Recovery rate	63%	> 75%	✗
Default rate	19%	< 15%	⚠
Death rate	0%	< 3%	✓
Referral rate	2%		

Water, Sanitation and Hygiene (WASH)

WATER

	INDICATOR	STANDARD
Average number of litres of potable water available per person per day		< 20
Proportion of households collecting >= 15 litres per person per day		> 95%
Number of persons of concern per water tap		< 80
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5		> 95%
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%
Proportion of households collecting drinking water from protected water sources only		> 95%
Proportion of households with sufficient daily water storage capacity		> 80%

WATER STAFFING

	INDICATOR	STANDARD
Number of hygiene promoters		1 : < 20,000

SANITATION

	INDICATOR	STANDARD
Refugees per latrine/toilet		< 20
Proportion of households with drop-hole latrine or drop-hole toilet		
Proportion of households report defecating in a toilet		> 85%
Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%