
Site report 2018 Panrieng SOUTH SUDAN

Population







Refugee Agency

Origin of refugees

Site opened May 2012

Partners



GAM



Key Indicators

Global Acute

CRUDE MORTALITY RATE

664.0

24.8 %

Malaria

Other

Deaths/1,000 refugees/

U5 MORTALITY RATE

Severe Acute

SAM

20.0 %

(with weapon) NC

Deaths/1,000 refugees/ month

MATERNAL MORTALITY

Deaths

WATER SUPPLY

Liters/ person/ day


SKILLED BIRTH ATTENDANCE


102% of all births

LATRINE COVERAGE


Refugees/communal

COMMUNICABLE DISEASES





NON-COMMUNICABLE DISEASES



Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARD
Number of medical doctors		1: < 50,000	•	Incidence of malaria among children under 5		
Number of qualified nurses		1:<10,000	•	Incidence of watery diarrhoea among		
Number of community health workers		1:<1,000	•	children under 5		
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5		
Consultations per trained clinician per day	49	< 50	②	Incidence of pneumonia among over 5	332	
Health utilisation rate (new visits / refugee /	306.0	1 - 4	•	Tuberculosis success rate		< 90%
year) Proportion of host population consultations	98%			Were any MDR/X-TB cases diagnosed among PoCs?		No
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	664.0	< 0.75	8	Full vaccination coverage		< 95%
Under-five Mortality Rate (U5MR) (/1000/month)		< 1.5		Measles vaccination coverage		< 95%
Infant Mortality Rate (IMR) (/1000 livebirths)	9.6	< 30	②	Source of vaccination data		
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	10	< 20	②	IN-PATIENT DEPARTMENT (IPD)	INDICATOR	STANDARD
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	22.8	
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0	
Number of outbreaks reported	0			Hospitalisation rate	110000.0	50 - 150

Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD
Antenatal care coverage	56%	> 90%	8	Contraceptive prevalence rate	6000%	≥ 30%
Coverage of antenatal tetanus vaccination	49%	> 95%	8	Source of contraceptive prevalence data		
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD
Proportion of births attended by skilled personnel	102%	≥ 90%	②	Total number of reported rapes	0	
Proportion of births conducted by caesarean section	69%	5 - 15%	②	Proportion of eligible rape survivors provided with PEP within 72 hours		100%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	4%	< 15%	②	Proportion of eligible rape survivors provided with ECP within 120 hours		100%
Proportion of teenage pregnancies	7%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD
Still birth rate (/ 1000 total births / month)	10			Number of genital ulcer diseases	1	
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	1	
Proportion of maternal deaths investigated within 48 hours	200%	100%	⊘			

HIV/AIDS

Death rate

Referral rate

PREVENTION & TREATMENT	INDICATOR	STANDARD		PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV	100%	100%	Ø	Number of PoCs receving ART	3	
Condom distribution rate	429.40	> 0.5	②	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%		PMTCT coverage	10%	100%
Nutrition						
NUTRITION GENERAL	INDICATOR	STANDARD		INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	Not collected			Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected	
Prevalence of stunting (6-59 months)	Not collected			Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected	
Prevalence of severe acute malnutrition (6 - 59 months)	Not	collected		IYCF programmes targeting pregnant and lactating women established or maintained?		Yes
Prevalence of anaemia in children (6-59 months)	Not	collected		MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not	collected		Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)		> 70%
CMAM: SAM TREATMENT	INDICATOR	STANDARD		MAM TREATMENT	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	0			Number of new moderate acute admissions (6-59 months)	0	
Recovery rate		> 75%		Recovery rate		> 75%
Default rate		< 0%		Default rate		< 15%

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD	WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day		< 20	Number of hygiene promoters		1:<20,000	•
Proportion of households collecting >= 15 litres per person per day		> 95%	SANITATION Refugees per latrine/toilet	INDICATOR	STANDARD < 20	
Number of persons of concern per water tap		< 80	Proportion of households with drop-hole		120	
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU		> 95%	latrine or drop-hole toilet Proportion of households report defecating in a toilet		> 85%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%	Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%	
Proportion of households collecting drinking water from protected water sources only		> 95%				
Proportion of households with sufficient daily water storage capacity		> 80%				

< 10%

Death rate

Referral rate

< 3%