
Site report 2018 Parmir SOUTH SUDAN

Population

27,794 refugees





Refugee Agency

Origin of refugees SUDAN

Site opened Sep 2016

Partners



Key Indicators

CRUDE MORTALITY RATE

Deaths/1,000 refugees/

6% Global Acute

GAM

U5 MORTALITY RATE

Deaths/1,000 refugees/

SAM

0% Severe Acute Malnutrition

MATERNAL MORTALITY

Deaths

WATER SUPPLY

25

Liters/ person/ day

SKILLED BIRTH **ATTENDANCE**

of all births


LATRINE COVERAGE


35.3 %

Refugees/communal toilets or latrines


12.9 %

COMMUNICABLE DISEASES





NON-COMMUNICABLE DISEASES



Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDAR	D	
Number of medical doctors		1:<50,000	•	Incidence of malaria among children under 5	38			
Number of qualified nurses		1 : < 10,000	•	Incidence of watery diarrhoea among	26			
Number of community health workers		1 : < 1,000	•	children under 5				
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	28			
Consultations per trained clinician per day	49	< 50	•	Incidence of pneumonia among over 5	11			
Health utilisation rate (new visits / refugee /	1.6	1 - 4	•	Tuberculosis success rate		< 90%		
year) Proportion of host population consultations	13%			Were any MDR/X-TB cases diagnosed among PoCs?	No			
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATOR	STAI	NDARD	
Crude Mortality Rate (CMR) (/1000/month)	0.1	< 0.75	②	Full vaccination coverage	74%	<	95%	8
Under-five Mortality Rate (U5MR) (/1000/month)	0.2	< 1.5	②	Measles vaccination coverage	74%	<	95%	8
Infant Mortality Rate (IMR) (/1000 livebirths)	5.0	< 30	②	Source of vaccination data				
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	3	< 20	②	IN-PATIENT DEPARTMENT (IPD)	INDICATOR	STAI	NDARD	
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	3.1			
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0			
Number of outbreaks reported	0			Hospitalisation rate	103.6	50	- 150	②

Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	87%	> 90%	<u> </u>	Contraceptive prevalence rate	1%	≥ 30%	8
Coverage of antenatal tetanus vaccination	93%	> 95%	<u> </u>	Source of contraceptive prevalence data			
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	93%	≥ 90%	②	Total number of reported rapes	11		
Proportion of births conducted by caesarean section	0%	5 - 15%	8	Proportion of eligible rape survivors provided with PEP within 72 hours	100%	100%	•
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	11%	< 15%	②	Proportion of eligible rape survivors provided with ECP within 120 hours	100%	100%	•
Proportion of teenage pregnancies	9%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate (/ 1000 total births / month)	1			Number of genital ulcer diseases	161		
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	155		
Proportion of maternal deaths investigated within 48 hours		100%					

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD		PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV	100%	100%	②	Number of PoCs receving ART	11	
Condom distribution rate	0.06	> 0.5	8	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%		PMTCT coverage	111%	100%

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD		INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD	
Prevalence of global acute malnutrition (6-59 months)	6%	< 20%	②	Proportion of children 0-23 months who were put to the breast within one hour of birth	84%	> 80%	8
Prevalence of stunting (6-59 months)	28%	< 20%	②	Proportion of infants aged between 0-5 months who received only breast milk during the previous day	88%	> 70%	8
Prevalence of severe acute malnutrition (6 - 59 months)	0%	< 3%	②	IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	
Prevalence of anaemia in children (6-59 months)	47%	< 20%	8	MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	23%	< 20%	A	Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)		> 70%	
CMAM: SAM TREATMENT	INDICATOR	STANDARD		MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)	114			Number of new moderate acute admissions (6-59 months)	713		
Recovery rate	86%	> 75%	②	Recovery rate	93%	> 75%	②
Default rate	7%	< 0%	8	Default rate	0%	< 15%	②
Death rate	0%	< 10%	②	Death rate	0%	< 3%	②
Referral rate	4%			Referral rate	6%		

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	25	< 20	②	Number of hygiene promoters		1:<20,000	•
Proportion of households collecting >= 15 litres per person per day		> 95%		SANITATION Refugees per latrine/toilet	INDICATOR 11	STANDARD < 20	o
Number of persons of concern per water tap		< 80		Proportion of households with drop-hole		- 20	
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5		> 95%		Proportion of households report defecating in a toilet		> 85%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%		Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%	
Proportion of households collecting drinking water from protected water sources only		> 95%					
Proportion of households with sufficient daily water storage capacity		> 80%					