



# Site report 2018 Yida SOUTH SUDAN



## Population

**51,530** refugees



Origin of refugees  
**SUDAN**

Site opened  
**Aug 2011**

## Partners



## Key Indicators

CRUDE MORTALITY RATE

**0.0** Deaths/1,000 refugees/month

U5 MORTALITY RATE

**0.0** Deaths/1,000 refugees/month

MATERNAL MORTALITY

**0** Deaths

SKILLED BIRTH ATTENDANCE

**81%** of all births

GAM

**NC** Global Acute Malnutrition

SAM

**NC** Severe Acute Malnutrition


WATER SUPPLY

Liters/ person/ day



LATRINE COVERAGE

Refugees/communal toilets or latrines


### COMMUNICABLE DISEASES




### INJURIES



### NON-COMMUNICABLE DISEASES



### MENTAL HEALTH



## Public Health

### HEALTH STAFFING

	INDICATOR	STANDARD
Number of medical doctors		1 : < 50,000
Number of qualified nurses		1 : < 10,000
Number of community health workers		1 : < 1,000

### ACCESS AND UTILIZATION

	INDICATOR	STANDARD
Consultations per trained clinician per day	99	< 50
Health utilisation rate (new visits / refugee / year)	1.3	1 - 4
Proportion of host population consultations	10%	

### MORTALITY

	INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.0	< 0.75
Under-five Mortality Rate (U5MR) (/1000/month)	0.0	< 1.5
Infant Mortality Rate (IMR) (/1000 livebirths)	0.6	< 30
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20

### OUTBREAK ALERT AND RESPONSE

	INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours		100%
Number of outbreaks reported	0	

### MORBIDITY

	INDICATOR	STANDARD
Incidence of malaria among children under 5	70	
Incidence of watery diarrhoea among children under 5	40	
Incidence of pneumonia among children under 5	4	
Incidence of pneumonia among over 5	2	
Tuberculosis success rate		< 90%
Were any MDR/X-TB cases diagnosed among PoCs?		No

### VACCINATION

	INDICATOR	STANDARD	
Full vaccination coverage	15%	< 95%	✗
Measles vaccination coverage	15%	< 95%	✗
Source of vaccination data			✓

### IN-PATIENT DEPARTMENT (IPD)

	INDICATOR	STANDARD	
Average length of stay (days)	0.1		
Case fatality rate	0.0		
Hospitalisation rate	8.7	50 - 150	✗

## Reproductive Health

### ANTENATAL CARE

	INDICATOR	STANDARD
Antenatal care coverage	37%	> 90%
Coverage of antenatal tetanus vaccination	94%	> 95%

### DELIVERY CARE

	INDICATOR	STANDARD
Proportion of births attended by skilled personnel	81%	≥ 90%
Proportion of births conducted by caesarean section	0%	5 - 15%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	1%	< 15%
Proportion of teenage pregnancies	9%	
Still birth rate ( / 1000 total births / month)	1	
Number of maternal deaths	0	
Proportion of maternal deaths investigated within 48 hours		100%

### FAMILY PLANNING

	INDICATOR	STANDARD	
Contraceptive prevalence rate	1%	≥ 30%	✗
Source of contraceptive prevalence data			⚠

### SEXUAL AND GENDER-BASED VIOLENCE (SGBV)

	INDICATOR	STANDARD	
Total number of reported rapes	8		⚠
Proportion of eligible rape survivors provided with PEP within 72 hours	80%	100%	✗
Proportion of eligible rape survivors provided with ECP within 120 hours	38%	100%	✗

### SEXUALLY TRANSMITTED INFECTIONS (STIS)

	INDICATOR	STANDARD
Number of genital ulcer diseases	11	
Number of Pelvic Inflammatory diseases	191	

## HIV/AIDS

### PREVENTION & TREATMENT

INDICATOR	STANDARD
Proportion of blood units screened for HIV	100%
Condom distribution rate	0.00 > 0.5
Proportion of TB patients tested for HIV	100%

### PREVENTION & TREATMENT

INDICATOR	STANDARD
Number of PoCs receiving ART	0
Proportion of HIV positive mothers who swallowed ARV during delivery	100%
PMTCT coverage	65% 100%

## Nutrition

### NUTRITION GENERAL

INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	Not collected
Prevalence of stunting (6-59 months)	Not collected
Prevalence of severe acute malnutrition (6 - 59 months)	Not collected
Prevalence of anaemia in children (6-59 months)	Not collected
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected

### INFANT AND YOUNG CHILD FEEDING (IYCF)

INDICATOR	STANDARD
Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected
Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected
IYCF programmes targeting pregnant and lactating women established or maintained?	Yes

### MICRONUTRIENT DEFICIENCY

INDICATOR	STANDARD
Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	> 70%

### CMAM: SAM TREATMENT

INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	631
Recovery rate	53% > 75%
Default rate	2% < 0%
Death rate	0% < 10%
Referral rate	42%

### MAM TREATMENT

INDICATOR	STANDARD
Number of new moderate acute admissions (6-59 months)	1600
Recovery rate	47% > 75%
Default rate	5% < 15%
Death rate	0% < 3%
Referral rate	47%

## Water, Sanitation and Hygiene (WASH)

### WATER

INDICATOR	STANDARD
Average number of litres of potable water available per person per day	< 20
Proportion of households collecting >= 15 litres per person per day	> 95%
Number of persons of concern per water tap	< 80
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual ( >= 0.1 mg/litre) and NTU <5	> 95%
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	> 95%
Proportion of households collecting drinking water from protected water sources only	> 95%
Proportion of households with sufficient daily water storage capacity	> 80%

### WATER STAFFING

INDICATOR	STANDARD
Number of hygiene promoters	1 : < 20,000

### SANITATION

INDICATOR	STANDARD
Refugees per latrine/toilet	< 20
Proportion of households with drop-hole latrine or drop-hole toilet	> 85%
Proportion of households report defecating in a toilet	> 90%
Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	> 90%