



# Site report 2018 Yusuf Batil SOUTH SUDAN



## Population

**44,592** refugees



### Origin of refugees

SUDAN NA

Site opened  
May 2012

## Partners



## Key Indicators

### CRUDE MORTALITY RATE

**0.1** Deaths/1,000 refugees/month

### U5 MORTALITY RATE

**0.3** Deaths/1,000 refugees/month

### MATERNAL MORTALITY

**3** Deaths

### SKILLED BIRTH ATTENDANCE

**79%** of all births

### GAM

**NC** Global Acute Malnutrition

### SAM

**NC** Severe Acute Malnutrition


### WATER SUPPLY

**21** Liters/ person/ day



### LATRINE COVERAGE

**13** Refugees/communal toilets or latrines


## COMMUNICABLE DISEASES




## INJURIES



## NON-COMMUNICABLE DISEASES



## MENTAL HEALTH



## Public Health

| HEALTH STAFFING                                       |           |              | MORBIDITY                   |  |                |
|---|-----------|--------------|-----------------------------|--|----------------|
|   | INDICATOR | STANDARD     |                             | INDICATOR  | STANDARD       |
| Number of medical doctors                             |           | 1 : < 50,000 | ⊖                           | Incidence of malaria among children under 5          | 12             |
| Number of qualified nurses                            |           | 1 : < 10,000 | ⊖                           | Incidence of watery diarrhoea among children under 5 | 64             |
| Number of community health workers                    |           | 1 : < 1,000  | ⊖                           | Incidence of pneumonia among children under 5        | 121            |
| ACCESS AND UTILIZATION                                |           |              | VACCINATION                 |  |                |
|   | INDICATOR | STANDARD     |                             | INDICATOR  | STANDARD       |
| Consultations per trained clinician per day           | 64        | < 50         | ⊗                           | Incidence of pneumonia among over 5                  | 44             |
| Health utilisation rate (new visits / refugee / year) | 2.8       | 1 - 4        | ✓                           | Tuberculosis success rate                            | < 90%          |
| Proportion of host population consultations           | 0%        |              |                             | Were any MDR/X-TB cases diagnosed among PoCs?        | No             |
| MORTALITY   |           |              | IN-PATIENT DEPARTMENT (IPD) |  |                |
|   | INDICATOR | STANDARD     |                             | INDICATOR  | STANDARD       |
| Crude Mortality Rate (CMR) (/1000/month)              | 0.1       | < 0.75       | ✓                           | Average length of stay (days)                        |                |
| Under-five Mortality Rate (U5MR) (/1000/month)        | 0.3       | < 1.5        | ✓                           | Case fatality rate                                   |                |
| Infant Mortality Rate (IMR) (/1000 livebirths)        | 9.8       | < 30         | ✓                           | Hospitalisation rate                                 | 0.0 50 - 150 ⊗ |
| Neonatal Mortality Rate (NNMR) (/1000 livebirths)     | 3         | < 20         | ✓                           |  |                |
| OUTBREAK ALERT AND RESPONSE                           |           |              |                             |  |                |
|   | INDICATOR | STANDARD     |                             |  |                |
| Proportion of outbreaks investigated within 48 hours  | 100%      | 100%         | ✓                           |  |                |
| Number of outbreaks reported                          | 13        |              |                             |  |                |

## Reproductive Health

| ANTENATAL CARE   |           |          | FAMILY PLANNING                         |  |             |
|--|-----------|----------|---|--|-------------|
|  | INDICATOR | STANDARD |   | INDICATOR  | STANDARD    |
| Antenatal care coverage  | 65%       | > 90%    | ⊗                                       | Contraceptive prevalence rate  | 10% ≥ 30% ⊗ |
| Coverage of antenatal tetanus vaccination  | 96%       | > 95%    | ✓                                       | Source of contraceptive prevalence data                                  |             |
| DELIVERY CARE  |           |          | SEXUAL AND GENDER-BASED VIOLENCE (SGBV) |  |             |
|  | INDICATOR | STANDARD |   | INDICATOR  | STANDARD    |
| Proportion of births attended by skilled personnel                                     | 79%       | ≥ 90%    | ⊗                                       | Total number of reported rapes   | 0           |
| Proportion of births conducted by caesarean section                                    | 0%        | 5 - 15%  | ⊗                                       | Proportion of eligible rape survivors provided with PEP within 72 hours  | 100%        |
| Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs) | 8%        | < 15%    | ✓                                       | Proportion of eligible rape survivors provided with ECP within 120 hours | 100%        |
| Proportion of teenage pregnancies  | 5%        |          |   | SEXUALLY TRANSMITTED INFECTIONS (STIS)                                   |             |
| Still birth rate ( / 1000 total births / month)  | 1         |          |   | Number of genital ulcer diseases   | 6           |
| Number of maternal deaths  | 3         |          |   | Number of Pelvic Inflammatory diseases                                   | 112         |
| Proportion of maternal deaths investigated within 48 hours                             | 100%      | 100%     | ✓                                       |  |             |

## HIV/AIDS

### PREVENTION & TREATMENT

Proportion of blood units screened for HIV

INDICATOR STANDARD

100%

Condom distribution rate

0.01 > 0.5

Proportion of TB patients tested for HIV

100%

### PREVENTION & TREATMENT

Number of PoCs receiving ART

INDICATOR STANDARD

4

✘ Proportion of HIV positive mothers who swallowed ARV during delivery

100%

PMTCT coverage

89% 100%



## Nutrition

### NUTRITION GENERAL

Prevalence of global acute malnutrition (6-59 months)

INDICATOR STANDARD

Not collected

Prevalence of stunting (6-59 months)

Not collected

Prevalence of severe acute malnutrition (6 - 59 months)

Not collected

Prevalence of anaemia in children (6-59 months)

Not collected

Prevalence of anaemia in women of reproductive age (15-49 yrs)

Not collected

### INFANT AND YOUNG CHILD FEEDING (IYCF)

Proportion of children 0-23 months who were put to the breast within one hour of birth

INDICATOR STANDARD

Not collected

Proportion of infants aged between 0-5 months who received only breast milk during the previous day

Not collected

IYCF programmes targeting pregnant and lactating women established or maintained?

Yes

### MICRONUTRIENT DEFICIENCY

Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)

INDICATOR STANDARD

> 70%

### CMAM: SAM TREATMENT

Number of new severe acute admissions (6-59 months)

INDICATOR STANDARD

156

Recovery rate

94% > 75%



Default rate

1% < 0%



Death rate

0% < 10%



Referral rate

1%

### MAM TREATMENT

Number of new moderate acute admissions (6-59 months)

INDICATOR STANDARD

833

Recovery rate

86% > 75%



Default rate

0% < 15%



Death rate

0% < 3%



Referral rate

7%

## Water, Sanitation and Hygiene (WASH)

### WATER

Average number of litres of potable water available per person per day

INDICATOR STANDARD

21 < 20



Proportion of households collecting >= 15 litres per person per day

> 95%

Number of persons of concern per water tap

< 80

Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual ( >= 0.1 mg/litre) and NTU <5

> 95%

Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml

> 95%

Proportion of households collecting drinking water from protected water sources only

> 95%

Proportion of households with sufficient daily water storage capacity

> 80%

### WATER STAFFING

Number of hygiene promoters

INDICATOR STANDARD

1 : < 20,000



### SANITATION

Refugees per latrine/toilet

INDICATOR STANDARD

13 < 20



Proportion of households with drop-hole latrine or drop-hole toilet

Proportion of households report defecating in a toilet

> 85%

Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)

> 90%