

# Site report 2018 Djabal CHAD



## Population


**22,577** refugees

**16%**

Children under 5

**20%**

Women of reproductive age



Origin of refugees

**SOUDAN**

Site opened

**Jan 2004**

## Partners



## Key Indicators

CRUDE MORTALITY RATE

**0.1** Deaths/1,000 refugees/month

U5 MORTALITY RATE

**0.5** Deaths/1,000 refugees/month

MATERNAL MORTALITY

**1** Deaths

SKILLED BIRTH ATTENDANCE

**98%** of all births

GAM

**NC** Global Acute Malnutrition

SAM

**NC** Severe Acute Malnutrition


WATER SUPPLY

**19** Liters/ person/ day



LATRINE COVERAGE

**6** Refugees/communal toilets or latrines


### COMMUNICABLE DISEASES




### INJURIES



### NON-COMMUNICABLE DISEASES



### MENTAL HEALTH



## Public Health

### HEALTH STAFFING

|                                    | INDICATOR | STANDARD     |
|------------------------------------|-----------|--------------|
| Number of medical doctors          | 1         | 1 : < 50,000 |
| Number of qualified nurses         | 4         | 1 : < 10,000 |
| Number of community health workers | 17        | 1 : < 1,000  |

### ACCESS AND UTILIZATION

|   | INDICATOR | STANDARD |
|---|-----------|----------|
| Consultations per trained clinician per day           | 16        | < 50     |
| Health utilisation rate (new visits / refugee / year) | 0.9       | 1 - 4    |
| Proportion of host population consultations           | 14%       |          |

### MORTALITY

|   | INDICATOR | STANDARD |
|---|-----------|----------|
| Crude Mortality Rate (CMR) (/1000/month)          | 0.1       | < 0.75   |
| Under-five Mortality Rate (U5MR) (/1000/month)    | 0.5       | < 1.5    |
| Infant Mortality Rate (IMR) (/1000 livebirths)    | 13.8      | < 30     |
| Neonatal Mortality Rate (NNMR) (/1000 livebirths) | 1         | < 20     |

### OUTBREAK ALERT AND RESPONSE

|  | INDICATOR | STANDARD |
|--|-----------|----------|
| Proportion of outbreaks investigated within 48 hours |           | 100%     |
| Number of outbreaks reported                         | 0         |          |

### MORBIDITY

|  | INDICATOR | STANDARD |
|--|-----------|----------|
| Incidence of malaria among children under 5          | 20        |          |
| Incidence of watery diarrhoea among children under 5 | 6         |          |
| Incidence of pneumonia among children under 5        | 86        |          |
| Incidence of pneumonia among over 5                  | 19        |          |
| Tuberculosis success rate                            | 100%      | < 90%    |
| Were any MDR/X-TB cases diagnosed among PoCs?        | No        | No       |

### VACCINATION

|                              | INDICATOR | STANDARD |
|------------------------------|-----------|----------|
| Full vaccination coverage    | 84%       | < 95%    |
| Measles vaccination coverage | 84%       | < 95%    |
| Source of vaccination data   | HIS       |          |

### IN-PATIENT DEPARTMENT (IPD)

|                               | INDICATOR | STANDARD |
|-------------------------------|-----------|----------|
| Average length of stay (days) | 3.4       |          |
| Case fatality rate            | 0.0       |          |
| Hospitalisation rate          | 14.6      | 50 - 150 |

## Reproductive Health

### ANTENATAL CARE

|   | INDICATOR | STANDARD |
|---|-----------|----------|
| Antenatal care coverage                   | 65%       | > 90%    |
| Coverage of antenatal tetanus vaccination | 97%       | > 95%    |

### DELIVERY CARE

|  | INDICATOR | STANDARD |
|--|-----------|----------|
| Proportion of births attended by skilled personnel                                     | 98%       | ≥ 90%    |
| Proportion of births conducted by caesarean section                                    | 1%        | 5 - 15%  |
| Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs) | 10%       | < 15%    |
| Proportion of teenage pregnancies  | 4%        |          |
| Still birth rate ( / 1000 total births / month)  | 1         |          |
| Number of maternal deaths  | 1         |          |
| Proportion of maternal deaths investigated within 48 hours                             | 100%      | 100%     |

### FAMILY PLANNING

|   | INDICATOR | STANDARD |
|---|-----------|----------|
| Contraceptive prevalence rate           | 42%       | ≥ 30%    |
| Source of contraceptive prevalence data | HIS       |          |

### SEXUAL AND GENDER-BASED VIOLENCE (SGBV)

|  | INDICATOR | STANDARD |
|--|-----------|----------|
| Total number of reported rapes   | 0         |          |
| Proportion of eligible rape survivors provided with PEP within 72 hours  |           | 100%     |
| Proportion of eligible rape survivors provided with ECP within 120 hours |           | 100%     |

### SEXUALLY TRANSMITTED INFECTIONS (STIS)

|  | INDICATOR | STANDARD |
|--|-----------|----------|
| Number of genital ulcer diseases       | 2         |          |
| Number of Pelvic Inflammatory diseases | 8         |          |

## HIV/AIDS

### PREVENTION & TREATMENT

| INDICATOR                                  | STANDARD   |
|--|------------|
| Proportion of blood units screened for HIV | 100%       |
| Condom distribution rate                   | 0.07 > 0.5 |
| Proportion of TB patients tested for HIV   | 100%       |

### PREVENTION & TREATMENT

| INDICATOR  | STANDARD |
|--|----------|
| Number of PoCs receiving ART   | 17       |
| Proportion of HIV positive mothers who swallowed ARV during delivery | 100%     |
| PMTCT coverage   | 90% 100% |

## Nutrition

### NUTRITION GENERAL

| INDICATOR  | STANDARD      |
|--|---------------|
| Prevalence of global acute malnutrition (6-59 months)          | Not collected |
| Prevalence of stunting (6-59 months)                           | Not collected |
| Prevalence of severe acute malnutrition (6 - 59 months)        | Not collected |
| Prevalence of anaemia in children (6-59 months)                | Not collected |
| Prevalence of anaemia in women of reproductive age (15-49 yrs) | Not collected |

### INFANT AND YOUNG CHILD FEEDING (IYCF)

| INDICATOR   | STANDARD      |
|---|---------------|
| Proportion of children 0-23 months who were put to the breast within one hour of birth              | Not collected |
| Proportion of infants aged between 0-5 months who received only breast milk during the previous day | Not collected |
| IYCF programmes targeting pregnant and lactating women established or maintained?                   | Yes           |

### MICRONUTRIENT DEFICIENCY

| INDICATOR   | STANDARD      |
|---|---------------|
| Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF) | Not collected |

### CMAM: SAM TREATMENT

| INDICATOR   | STANDARD  |
|---|-----------|
| Number of new severe acute admissions (6-59 months) | 360       |
| Recovery rate                                       | 81% > 75% |
| Default rate  | 4% < 0%   |
| Death rate  | 0% < 10%  |
| Referral rate                                       | 14%       |

### MAM TREATMENT

| INDICATOR   | STANDARD  |
|---|-----------|
| Number of new moderate acute admissions (6-59 months) | 634       |
| Recovery rate   | 82% > 75% |
| Default rate  | 5% < 15%  |
| Death rate  | 0% < 3%   |
| Referral rate   | 12%       |

## Water, Sanitation and Hygiene (WASH)

### WATER

| INDICATOR  | STANDARD   |
|--|------------|
| Average number of litres of potable water available per person per day   | 19 < 20    |
| Proportion of households collecting >= 15 litres per person per day  | 100% > 95% |
| Number of persons of concern per water tap   | 68 < 80    |
| Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual ( >= 0.1 mg/litre) and NTU <5 | 100% > 95% |
| Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml                                    | 100% > 95% |
| Proportion of households collecting drinking water from protected water sources only   | 100% > 95% |
| Proportion of households with sufficient daily water storage capacity  | 100% > 80% |

### WATER STAFFING

| INDICATOR                   | STANDARD       |
|-----------------------------|----------------|
| Number of hygiene promoters | 2 1 : < 20,000 |

### SANITATION

| INDICATOR   | STANDARD   |
|---|------------|
| Refugees per latrine/toilet   | 6 < 20     |
| Proportion of households with drop-hole latrine or drop-hole toilet   | 44%        |
| Proportion of households report defecating in a toilet  | 44% > 85%  |
| Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety) | 100% > 90% |