

Site report 2018 Moyo CHAD



Population

8,397 refugees

16%

Children under 5

20%

Women of reproductive age



Origin of refugees

CENTRAFRIQUE

Site opened

Apr 2011

Partners



Key Indicators

CRUDE MORTALITY RATE

0.1 Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.4 Deaths/1,000 refugees/month

MATERNAL MORTALITY

0 Deaths

SKILLED BIRTH ATTENDANCE

100% of all births

GAM

NC Global Acute Malnutrition

SAM

NC Severe Acute Malnutrition


WATER SUPPLY

38 Liters/ person/ day


LATRINE COVERAGE

6 Refugees/communal toilets or latrines


COMMUNICABLE DISEASES




INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING			MORBIDITY			
	INDICATOR	STANDARD		INDICATOR	STANDARD	
Number of medical doctors	1	1 : < 50,000	✓	90		
Number of qualified nurses	3	1 : < 10,000	✓	6		
Number of community health workers	19	1 : < 1,000	✓	120		
ACCESS AND UTILIZATION			VACCINATION			
	INDICATOR	STANDARD		INDICATOR	STANDARD	
Consultations per trained clinician per day	45	< 50	✓	91%	< 95% 	
Health utilisation rate (new visits / refugee / year)	1.6	1 - 4	✓	Measles vaccination coverage	Not collected	
Proportion of host population consultations	48%		✓	Source of vaccination data	HIS	
MORTALITY			IN-PATIENT DEPARTMENT (IPD)			
	INDICATOR	STANDARD		INDICATOR	STANDARD	
Crude Mortality Rate (CMR) (/1000/month)	0.1	< 0.75	✓	Average length of stay (days)	5.4	
Under-five Mortality Rate (U5MR) (/1000/month)	0.4	< 1.5	✓	Case fatality rate	0.0	
Infant Mortality Rate (IMR) (/1000 livebirths)	14.3	< 30	✓	Hospitalisation rate	32.3	50 - 150 
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	✓	OUTBREAK ALERT AND RESPONSE		
	INDICATOR	STANDARD		INDICATOR	STANDARD	
Proportion of outbreaks investigated within 48 hours		100%		Proportion of outbreaks investigated within 48 hours		
Number of outbreaks reported	0			Number of outbreaks reported		

Reproductive Health

ANTENATAL CARE			FAMILY PLANNING			
	INDICATOR	STANDARD		INDICATOR	STANDARD	
Antenatal care coverage	91%	> 90%	✓	33%	≥ 30% 	
Coverage of antenatal tetanus vaccination	97%	> 95%	✓	Source of contraceptive prevalence data	HIS	
DELIVERY CARE			SEXUAL AND GENDER-BASED VIOLENCE (SGBV)			
	INDICATOR	STANDARD		INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	100%	≥ 90%	✓	Total number of reported rapes	0	
Proportion of births conducted by caesarean section	4%	5 - 15%	✗	Proportion of eligible rape survivors provided with PEP within 72 hours	100%	
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	3%	< 15%	✓	Proportion of eligible rape survivors provided with ECP within 120 hours	100%	
Proportion of teenage pregnancies	10%			SEXUALLY TRANSMITTED INFECTIONS (STIS)		
Still birth rate (/ 1000 total births / month)	3			Number of genital ulcer diseases	16	
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	19	
Proportion of maternal deaths investigated within 48 hours		100%				

HIV/AIDS

PREVENTION & TREATMENT

Proportion of blood units screened for HIV

100%

100%



Number of PoCs receiving ART

45

Condom distribution rate

0.20

> 0.5



Proportion of HIV positive mothers who swallowed ARV during delivery

100%

Proportion of TB patients tested for HIV

100%

PMTCT coverage

100%

100%



Nutrition

NUTRITION GENERAL

Prevalence of global acute malnutrition (6-59 months)

Not collected

Prevalence of stunting (6-59 months)

Not collected

Prevalence of severe acute malnutrition (6 - 59 months)

Not collected

Prevalence of anaemia in children (6-59 months)

Not collected

Prevalence of anaemia in women of reproductive age (15-49 yrs)

Not collected

INFANT AND YOUNG CHILD FEEDING (IYCF)

Proportion of children 0-23 months who were put to the breast within one hour of birth

Not collected

Proportion of infants aged between 0-5 months who received only breast milk during the previous day

Not collected

IYCF programmes targeting pregnant and lactating women established or maintained?

Yes

MICRONUTRIENT DEFICIENCY

Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)

56%

> 70%



CMAM: SAM TREATMENT

Number of new severe acute admissions (6-59 months)

95

Recovery rate

98%

> 75%



Default rate

0%

< 0%



Death rate

3%

< 10%



Referral rate

0%

MAM TREATMENT

Number of new moderate acute admissions (6-59 months)

99

Recovery rate

100%

> 75%



Default rate

0%

< 15%



Death rate

0%

< 3%



Referral rate

0%

Water, Sanitation and Hygiene (WASH)

WATER

Average number of litres of potable water available per person per day

38

< 20



Proportion of households collecting >= 15 litres per person per day

100%

> 95%



Number of persons of concern per water tap

381

< 80



Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5

0%

> 95%



Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml

100%

> 95%



Proportion of households collecting drinking water from protected water sources only

100%

> 95%



Proportion of households with sufficient daily water storage capacity

100%

> 80%



WATER STAFFING

Number of hygiene promoters

7

1 : < 20,000



SANITATION

Refugees per latrine/toilet

6

< 20



Proportion of households with drop-hole latrine or drop-hole toilet

61%

Proportion of households report defecating in a toilet

61%

> 85%



Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)

100%

> 90%

