

Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARD		
Number of medical doctors	1	1 : < 50,000	0	Incidence of malaria among children under 5	63			
Number of qualified nurses	3	1 : < 10,000	0	Incidence of watery diarrhoea among	10			
Number of community health workers	8	1 : < 1,000	8	children under 5				
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	27			
Consultations per trained clinician per day	49	< 50	0	Incidence of pneumonia among over 5	7			
Health utilisation rate (new visits / refugee /	1.2	1 - 4	0	Tuberculosis success rate	100%	< 90%	0	
year) Proportion of host population consultations	33%			Were any MDR/X-TB cases diagnosed among PoCs?	No	No	8	
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATO	STAND	ARD	
Crude Mortality Rate (CMR) (/1000/month)	0.2	< 0.75	0	Full vaccination coverage	89%	< 95	%	▲
Under-five Mortality Rate (U5MR) (/1000/month)	0.4	< 1.5	0	Measles vaccination coverage		Not collected		
Infant Mortality Rate (IMR) (/1000 livebirths)	5.4	< 30	0	Source of vaccination data	HIS			
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	5	< 20	0	IN-PATIENT DEPARTMENT (IPD)	INDICATO	STAND.	ARD	
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	1.2			
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0			
Number of outbreaks reported	0			Hospitalisation rate	121.9	50 - 1	50	0

Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	94%	> 90%	0	Contraceptive prevalence rate	95%	≥ 30%	0
Coverage of antenatal tetanus vaccination	95%	> 95%	4	Source of contraceptive prevalence data	HIS		
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	95%	≥ 90%	0	Total number of reported rapes	6		
Proportion of births conducted by caesarean section	3%	5 - 15%	8	Proportion of eligible rape survivors provided with PEP within 72 hours	160%	100%	O
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	4%	< 15%	0	Proportion of eligible rape survivors provided with ECP within 120 hours	50%	100%	8
Proportion of teenage pregnancies	9%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate (/ 1000 total births / month)	3			Number of genital ulcer diseases	46		
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	50		
Proportion of maternal deaths investigated within 48 hours		100%					

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD		PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV	100%	100%	0	Number of PoCs receving ART	151	
Condom distribution rate	0.12	> 0.5	8	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%		PMTCT coverage	99%	100%

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD		INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD	
Prevalence of global acute malnutrition (6-59 months)	Not collected			Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected		
Prevalence of stunting (6-59 months)	Not collected			Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected		
Prevalence of severe acute malnutrition (6 - 59 months)	Not collected			IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	
Prevalence of anaemia in children (6-59 months)	Not collected			MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected			Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	40%	> 70%	⊗
CMAM: SAM TREATMENT	INDICATOR	STANDARD		MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)	130			Number of new moderate acute admissions (6-59 months)	284		
Recovery rate	98%	> 75%	0	Recovery rate	91%	> 75%	0
Default rate	0%	< 0%	8	Default rate	0%	< 15%	⊘
Death rate	0%	< 10%	0	Death rate	0%	< 3%	⊘
Referral rate	0%			Referral rate	8%		

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	46	< 20	0	Number of hygiene promoters	7	1 : < 20,000	0
Proportion of households collecting >= 15 litres per person per day	100%	> 95%	0	SANITATION	INDICATOR	STANDARD	0
Number of persons of concern per water tap	530	< 80	8	Refugees per latrine/toilet Proportion of households with drop-hole	65%	< 20	
Proportion of tests at chlorinated water	100%	> 95%	0	latrine or drop-hole toilet			•
collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5				Proportion of households report defecating in a toilet	65%	> 85%	8
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	97%	> 95%	0	Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	100%	> 90%	0
Proportion of households collecting drinking water from protected water sources only	100%	> 95%	0				
Proportion of households with sufficient daily water storage capacity	100%	> 80%	0				