

, intellectual disability

# **Public Health**

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARD	
Number of medical doctors	1	1 : < 50,000	▲	Incidence of malaria among children under 5	3		
Number of qualified nurses	7	1 : < 10,000	0	Incidence of watery diarrhoea among	20		
Number of community health workers	37	1 : < 1,000	۲	children under 5			
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	19		
Consultations per trained clinician per day	28	< 50	0	Incidence of pneumonia among over 5	4		
Health utilisation rate (new visits / refugee /	0.7	1 - 4	8	Tuberculosis success rate	100%	< 90%	0
year) Proportion of host population consultations	11%			Were any MDR/X-TB cases diagnosed among PoCs?	No	No	8
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATOR	STANDA	RD
Crude Mortality Rate (CMR) (/1000/month)	0.2	< 0.75	0	Full vaccination coverage	93%	< 95%	4
Under-five Mortality Rate (U5MR) (/1000/month)	0.5	< 1.5	0	Measles vaccination coverage	96%	< 95%	0
Infant Mortality Rate (IMR) (/1000 livebirths)	22.2	< 30	0	Source of vaccination data	HIS		
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	0	IN-PATIENT DEPARTMENT (IPD)	INDICATOR	STANDA	RD
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	2.0		
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0		
Number of outbreaks reported	0			Hospitalisation rate	18.2	50 - 15	0 😣

### Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	97%	> 90%	0	Contraceptive prevalence rate	52%	≥ 30%	<b>⊘</b>
Coverage of antenatal tetanus vaccination	98%	> 95%	<b>~</b>	Source of contraceptive prevalence data	HIS		
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	100%	≥ 90%	0	Total number of reported rapes	0		
Proportion of births conducted by caesarean section	0%	5 - 15%	⊗	Proportion of eligible rape survivors provided with PEP within 72 hours		100%	
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	1%	< 15%	0	Proportion of eligible rape survivors provided with ECP within 120 hours		100%	
Proportion of teenage pregnancies	3%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate ( / 1000 total births / month)	1			Number of genital ulcer diseases	13		
Number of maternal deaths	1			Number of Pelvic Inflammatory diseases	23		
Proportion of maternal deaths investigated within 48 hours	100%	100%	0				

#### HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD		PREVENTION & TREATMENT	INDICATOR	STANDARD	
Proportion of blood units screened for HIV	100%	100%	0	Number of PoCs receving ART	1		
Condom distribution rate	0.01	> 0.5	8	Proportion of HIV positive mothers who swallowed ARV during delivery		100%	
Proportion of TB patients tested for HIV		100%		PMTCT coverage	89%	100%	8

# Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD		INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD	
Prevalence of global acute malnutrition (6-59 months)	Not collected			Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected		
Prevalence of stunting (6-59 months)	Not collected			Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected		
Prevalence of severe acute malnutrition (6 - 59 months)	Not collected			IYCF programmes targeting pregnant and lactating women established or maintained?	Yes		
Prevalence of anaemia in children (6-59 months)	Not collected			MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected			Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	Not collected		
CMAM: SAM TREATMENT	INDICATOR	STANDARD		MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)	807			Number of new moderate acute admissions (6-59 months)	1096		
Recovery rate	95%	> 75%	0	Recovery rate	81%	> 75%	<b>I</b>
Default rate	1%	< 0%	8	Default rate	0%	< 15%	<b>⊘</b>
Death rate	2%	< 10%	0	Death rate	0%	< 3%	0
Referral rate	2%			Referral rate	19%		

# Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	14	< 20	8	Number of hygiene promoters	14	1 : < 20,000	0
Proportion of households collecting >= 15 litres per person per day	74%	> 95%	▲	SANITATION	INDICATOR	STANDARD	0
Number of persons of concern per water tap	148	< 80	8	Refugees per latrine/toilet Proportion of households with drop-hole	9 47%	< 20	
Proportion of tests at chlorinated water	96%	> 95%	0	latrine or drop-hole toilet			
collection points indicating a measurable chlorine residual ( >= 0.1 mg/litre) and NTU <5				Proportion of households report defecating in a toilet	93%	> 85%	<b>S</b>
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	0%	> 95%	8	Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	0%	> 90%	8
Proportion of households collecting drinking water from protected water sources only	98%	> 95%	0				
Proportion of households with sufficient daily water storage capacity	100%	> 80%	0				