





Origin of refugees


Site opened Jan 2015

Partners




Key Indicators


SKILLED BIRTH CRUDE MORTALITY RATE **U5 MORTALITY RATE** MATERNAL MORTALITY ATTENDANCE Deaths/1,000 refugees/ 80% 0.1Deaths/1,000 refugees/ 0.30 of all births Deaths month month SAM WATER SUPPLY LATRINE COVERAGE GAM NC Global Acute NC Severe Acute 24 Refugees/communal 18 Liters/ person/ day Malnutrition Malnutrition toilets or latrines COMMUNICABLE DISEASES **NON-COMMUNICABLE DISEASES** 14.5 % 55.1 % Malaria Cardiovascular 11.8 % (confirm disease Malaria 16.0 % (suspected) LRTI 10.9 % 13.8 % 11.2 % Intestinal Malaria Malaria worms



74.5 %

Other





use disorder

Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARD		
Number of medical doctors	1	1 : < 50,000	0	Incidence of malaria among children under 5	26			
Number of qualified nurses	4	1 : < 10,000	0	Incidence of watery diarrhoea among	27			
Number of community health workers	33	1 : < 1,000	0	children under 5				
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	58			
Consultations per trained clinician per day	25	< 50	0	Incidence of pneumonia among over 5	15			
Health utilisation rate (new visits / refugee /	1.1	1 - 4	0	Tuberculosis success rate	100%	< 90%	0	
year) Proportion of host population consultations	52%			Were any MDR/X-TB cases diagnosed among PoCs?	No	No	8	
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATO	STAND	OARD	
Crude Mortality Rate (CMR) (/1000/month)	0.1	< 0.75	0	Full vaccination coverage	73%	< 95	5%	⊗
Under-five Mortality Rate (U5MR) (/1000/month)	0.3	< 1.5	0	Measles vaccination coverage	Not collected			
Infant Mortality Rate (IMR) (/1000 livebirths)	6.7	< 30	0	Source of vaccination data	HIS			
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	0	IN-PATIENT DEPARTMENT (IPD)	INDICATO	RSTAND	OARD	
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	0.1			
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0			
Number of outbreaks reported	0			Hospitalisation rate	36.7	50 - 1	150	8

Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	78%	> 90%	8	Contraceptive prevalence rate	415%	≥ 30%	⊘
Coverage of antenatal tetanus vaccination	96%	> 95%	0	Source of contraceptive prevalence data	HIS		
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	80%	≥ 90%	4	Total number of reported rapes	11		
Proportion of births conducted by caesarean section	1%	5 - 15%	⊗	Proportion of eligible rape survivors provided with PEP within 72 hours	0%	100%	8
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	3%	< 15%	0	Proportion of eligible rape survivors provided with ECP within 120 hours	0%	100%	⊗
Proportion of teenage pregnancies	15%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate (/ 1000 total births / month)	3			Number of genital ulcer diseases	0		
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	71		
Proportion of maternal deaths investigated within 48 hours		100%					

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD	PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%	Number of PoCs receving ART	217	
Condom distribution rate	0.17	> 0.5	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%	PMTCT coverage	110%	100% 📀

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD		INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD	
Prevalence of global acute malnutrition (6-59 months)	Not collected			Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected		
Prevalence of stunting (6-59 months)	Not collected			Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected		
Prevalence of severe acute malnutrition (6 - 59 months)	Not collected			IYCF programmes targeting pregnant and lactating women established or maintained?	Yes		
Prevalence of anaemia in children (6-59 months)	Not collected			MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected			Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	56%	> 70%	▲
CMAM: SAM TREATMENT	INDICATOR	STANDARD		MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)	158			Number of new moderate acute admissions (6-59 months)	411		
Recovery rate	94%	> 75%	0	Recovery rate	91%	> 75%	⊘
Default rate	1%	< 0%	8	Default rate	4%	< 15%	0
Death rate	1%	< 10%	0	Death rate	0%	< 3%	0
Referral rate	5%			Referral rate	3%		

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	24	< 20	0	Number of hygiene promoters	12	1 : < 20,000	0
Proportion of households collecting >= 15 litres per person per day	100%	> 95%	0	SANITATION Refugees per latrine/toilet	INDICATOR	STANDARD	•
Number of persons of concern per water tap	696	< 80	8	Proportion of households with drop-hole latrine or drop-hole toilet	14%		
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5	0%	> 95%	8	Proportion of households report defecating in a toilet	99%	> 85%	0
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	100%	> 95%	0	Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	39%	> 90%	8
Proportion of households collecting drinking water from protected water sources only	100%	> 95%	0				
Proportion of households with sufficient daily water storage capacity	99%	> 80%	0				