
# Site report 2018 Dosseye CHAD

## **Population**

16,081 refugees







# Origin of refugees CENTRAFRIQUE

Site opened Jan 2006

#### **Partners**





## **Key Indicators**

**CRUDE MORTALITY RATE** 

0.2

Deaths/1,000 refugees/

GAM

7% Global Acute

U5 MORTALITY RATE

0.7

Deaths/1,000 refugees/

SAM

1%

Severe Acute Malnutrition MATERNAL MORTALITY

0

Deaths

WATER SUPPLY

40

Liters/ person/ day

SKILLED BIRTH ATTENDANCE

97%


of all births

LATRINE COVERAGE


6


Refugees/communal toilets or latrines

#### COMMUNICABLE DISEASES




#### **INJURIES**





#### **NON-COMMUNICABLE DISEASES**



## **Public Health**

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARD		
Number of medical doctors	1	1: < 50,000	•	Incidence of malaria among children under 5	92			
Number of qualified nurses	4	1:<10,000	<b>②</b>	Incidence of watery diarrhoea among	4			
Number of community health workers	8	1:<1,000	8	children under 5				
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	77			
Consultations per trained clinician per day	27	< 50	<b>②</b>	Incidence of pneumonia among over 5	18			
Health utilisation rate (new visits / refugee /	1.3	1 - 4	•	Tuberculosis success rate	100%	< 90%	<b>②</b>	
year)  Proportion of host population consultations	14%			Were any MDR/X-TB cases diagnosed among PoCs?	No	No	8	
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATOR	STANDA	ARD	
Crude Mortality Rate (CMR) (/1000/month)	0.2	< 0.75	<b>②</b>	Full vaccination coverage	88%	< 959	%	<u> </u>
Under-five Mortality Rate (U5MR) (/1000/month)	0.7	< 1.5	<b>②</b>	Measles vaccination coverage	80%	< 959	%	<u> </u>
Infant Mortality Rate (IMR) (/1000 livebirths)	25.4	< 30	<b>②</b>	Source of vaccination data	HIS			
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	<b>②</b>	IN-PATIENT DEPARTMENT (IPD)	INDICATOR	STANDA	ARD	
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	2.3			
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0			
Number of outbreaks reported	0			Hospitalisation rate	94.6	50 - 1	50	<b>②</b>

## Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	98%	> 90%	<b>②</b>	Contraceptive prevalence rate	56%	≥ 30%	
Coverage of antenatal tetanus vaccination	98%	> 95%	<b>②</b>	Source of contraceptive prevalence data	HIS		
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	97%	≥ 90%	<b>②</b>	Total number of reported rapes	0		
Proportion of births conducted by caesarean section	0%	5 - 15%	8	Proportion of eligible rape survivors provided with PEP within 72 hours		100%	
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	1%	< 15%	<b>②</b>	Proportion of eligible rape survivors provided with ECP within 120 hours		100%	
Proportion of teenage pregnancies	17%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate ( / 1000 total births / month)	1			Number of genital ulcer diseases	35		
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	44		
Proportion of maternal deaths investigated within 48 hours		100%					

### HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD		PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV	100%	100%	<b>②</b>	Number of PoCs receving ART	139	
Condom distribution rate	0.12	> 0.5	8	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%		PMTCT coverage	100%	100%

## **Nutrition**

NUTRITION GENERAL	INDICATOR	STANDARD		INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD	
Prevalence of global acute malnutrition (6-59 months)	7% < 20%		<b>②</b>	Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected		
Prevalence of stunting (6-59 months)	42% < 20%		<b>②</b>	Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected		
Prevalence of severe acute malnutrition (6 - 59 months)	1%	< 3%	<b>②</b>	IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	
Prevalence of anaemia in children (6-59 months)	38%	< 20%	8	MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected			Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	62%	> 70%	<b>A</b>
CMAM: SAM TREATMENT	INDICATOR	STANDARD		MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)	179			Number of new moderate acute admissions (6-59 months)	253		
Recovery rate	94%	> 75%	<b>②</b>	Recovery rate	95%	> 75%	<b>②</b>
Default rate	1%	< 0%	8	Default rate	0%	< 15%	<b>②</b>
Death rate	0%	< 10%	<b>②</b>	Death rate	0%	< 3%	<b>②</b>
Referral rate	0%			Referral rate	5%		

## Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD		
Average number of litres of potable water available per person per day	40	< 20	<b>②</b>	Number of hygiene promoters	5	1:<20,000	<b>Ø</b>	
Proportion of households collecting >= 15 litres per person per day	100%	> 95%	•	SANITATION  Refugees per latrine/toilet	INDICATOR 6	STANDARD < 20	<b>O</b>	
Number of persons of concern per water tap	458	< 80	8	Proportion of households with drop-hole latrine or drop-hole toilet	64%			
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual ( >= 0.1 mg/litre) and NTU <5	0%	> 95%	8	Proportion of households report defecating in a toilet	64%	> 85%	8	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	100%	> 95%	•	Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	100%	> 90%	•	
Proportion of households collecting drinking water from protected water sources only	100%	> 95%	•					
Proportion of households with sufficient daily water storage capacity	100%	> 80%	•					