
Site report 2018 Gaga CHAD

Population

25,816 refugees







Origin of refugees SOUDAN

Site opened Jan 2005

Partners



Key Indicators

CRUDE MORTALITY RATE

Deaths/1,000 refugees/

month

GAM

Global Acute

U5 MORTALITY RATE

Deaths/1,000 refugees/ month

SAM

NC Severe Acute MATERNAL MORTALITY

Deaths

WATER SUPPLY

Liters/ person/ day


SKILLED BIRTH ATTENDANCE


100% of all births

LATRINE COVERAGE


Refugees/communal toilets or latrines

COMMUNICABLE DISEASES





NON-COMMUNICABLE DISEASES



Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARD		
Number of medical doctors	0	1: < 50,000	8	Incidence of malaria among children under 5	3			
Number of qualified nurses	6	1:<10,000	②	Incidence of watery diarrhoea among	20			
Number of community health workers	17	1:<1,000	8	children under 5				
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	27			
Consultations per trained clinician per day	31	< 50	•	Incidence of pneumonia among over 5	7			
Health utilisation rate (new visits / refugee /	0.7	1 - 4	8	Tuberculosis success rate	100%	< 90%	•	
year) Proportion of host population consultations	34%			Were any MDR/X-TB cases diagnosed among PoCs?	No	No	8	
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATO	R	ARD	
Crude Mortality Rate (CMR) (/1000/month)	0.3	< 0.75	②	Full vaccination coverage	99%	< 95	%	•
Under-five Mortality Rate (U5MR) (/1000/month)	0.7	< 1.5	②	Measles vaccination coverage	99%	< 95	%	•
Infant Mortality Rate (IMR) (/1000 livebirths)	30.2	< 30	Δ	Source of vaccination data	HIS			
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	②	IN-PATIENT DEPARTMENT (IPD)	INDICATO	R STAND	ARD	
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	2.6			
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0			
Number of outbreaks reported	0			Hospitalisation rate	8.1	50 - 1	50	•

Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	93%	> 90%	②	Contraceptive prevalence rate	57%	≥ 30%	•
Coverage of antenatal tetanus vaccination	95%	> 95%	<u> </u>	Source of contraceptive prevalence data	HIS		
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	100%	≥ 90%	②	Total number of reported rapes	1		
Proportion of births conducted by caesarean section	0%	5 - 15%	8	Proportion of eligible rape survivors provided with PEP within 72 hours		100%	
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	2%	< 15%	②	Proportion of eligible rape survivors provided with ECP within 120 hours	100%	100%	•
Proportion of teenage pregnancies	5%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate (/ 1000 total births / month)	1			Number of genital ulcer diseases	7		
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	22		
Proportion of maternal deaths investigated within 48 hours	200%	100%	②				

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD		PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%		Number of PoCs receving ART	13	
Condom distribution rate	0.01	> 0.5	8	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%		PMTCT coverage	100%	100%

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD		INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD	
Prevalence of global acute malnutrition (6-59 months)	Not collected			Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected		
Prevalence of stunting (6-59 months)	Not collected			Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected		
Prevalence of severe acute malnutrition (6 - 59 months)	Not collected			IYCF programmes targeting pregnant and lactating women established or maintained?	Yes		
Prevalence of anaemia in children (6-59 months)	Not collected			MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected			Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	Not collected		
CMAM: SAM TREATMENT	INDICATOR	STANDARD		MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)	479			Number of new moderate acute admissions (6-59 months)	616		
Recovery rate	79%	> 75%	②	Recovery rate	86%	> 75%	②
Default rate	5%	< 0%	8	Default rate	2%	< 15%	②
Death rate	1%	< 10%	②	Death rate	0%	< 3%	②
Referral rate	15%			Referral rate	12%		

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	9	< 20	8	Number of hygiene promoters	7	1:<20,000	•
Proportion of households collecting >= 15 litres per person per day	100%	> 95%	②	SANITATION Refugees per latrine/toilet	INDICATOR 7	STANDARD < 20	•
Number of persons of concern per water tap	115	< 80	8	Proportion of households with drop-hole	61%	- 20	
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5	93%	> 95%	A	Proportion of households report defecating in a toilet	92%	> 85%	•
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	0%	> 95%	8	Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	100%	> 90%	•
Proportion of households collecting drinking water from protected water sources only	59%	> 95%	8				
Proportion of households with sufficient daily water storage capacity	100%	> 80%	•				