Site report 2018 Gondje CHAD

Population

9,029 refugees







Origin of refugees CENTRAFRIQUE

Site opened Jan 2005

Partners





Key Indicators

CRUDE MORTALITY RATE

Deaths/1,000 refugees/

GAM

NC Global Acute U5 MORTALITY RATE

Deaths/1,000 refugees/

SAM

NC

Severe Acute Malnutrition

MATERNAL MORTALITY

0

Deaths

WATER SUPPLY

32

Liters/ person/ day

SKILLED BIRTH **ATTENDANCE**


91% of all births


LATRINE COVERAGE


6

Refugees/communal toilets or latrines


COMMUNICABLE DISEASES








NON-COMMUNICABLE DISEASES





Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARI	O .	
Number of medical doctors	1	1:<50,000	•	Incidence of malaria among children under 5	112			
Number of qualified nurses	2	1:<10,000	②	Incidence of watery diarrhoea among	16			
Number of community health workers	7	1:<1,000	8	children under 5				
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	151			
Consultations per trained clinician per day	51	< 50	A	Incidence of pneumonia among over 5	38			
Health utilisation rate (new visits / refugee /	2.5	1 - 4	•	Tuberculosis success rate	100%	< 90%	Ø	
year)				Were any MDR/X-TB cases diagnosed among	No	No	8	
Proportion of host population consultations	29%			PoCs?				
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATO	STAN	IDARD	
Crude Mortality Rate (CMR) (/1000/month)	0.1	< 0.75	⊘	Full vaccination coverage	91%	< !	95%	4
Under-five Mortality Rate (U5MR) (/1000/month)	0.6	< 1.5	②	Measles vaccination coverage		Not collecte	ed	
Infant Mortality Rate (IMR) (/1000 livebirths)	5.6	< 30	②	Source of vaccination data	HIS			
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	②	IN-PATIENT DEPARTMENT (IPD)	INDICATO	STAN	IDARD	
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	1.4			
Proportion of outbreaks investigated within 4 hours	8	100%		Case fatality rate	0.0			
Number of outbreaks reported	0			Hospitalisation rate	76.4	50	- 150	•

Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	74%	> 90%	8	Contraceptive prevalence rate	79%	≥ 30%	
Coverage of antenatal tetanus vaccination	93%	> 95%	<u> </u>	Source of contraceptive prevalence data	HIS		
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	91%	≥ 90%	②	Total number of reported rapes	0		
Proportion of births conducted by caesarean section	0%	5 - 15%	8	Proportion of eligible rape survivors provided with PEP within 72 hours		100%	
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	9%	< 15%	②	Proportion of eligible rape survivors provided with ECP within 120 hours		100%	
Proportion of teenage pregnancies	10%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate (/ 1000 total births / month)	0			Number of genital ulcer diseases	117		
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	75		
Proportion of maternal deaths investigated within 48 hours		100%					

HIV/AIDS

PREVENTION & TREATMENT

Condom distribution rate

Death rate

Referral rate

Proportion of blood units screened for HIV

			8	swallowed ARV during delivery			
Proportion of TB patients tested for HIV		100%		PMTCT coverage	98%	100%	A
Nutrition							
NUTRITION GENERAL	INDICATOR	STANDARD		INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD	
Prevalence of global acute malnutrition (6-59 months)	Not collected			Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected		
Prevalence of stunting (6-59 months)	Not	collected		Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not	collected	
Prevalence of severe acute malnutrition (6 - 59 months)	Not collected			IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	
Prevalence of anaemia in children (6-59 months)	Not collected			MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected			Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	48%	> 70%	8
CMAM: SAM TREATMENT	INDICATOR	STANDARD		MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)	137			Number of new moderate acute admissions (6-59 months)	249		
Recovery rate	100%	> 75%	②	Recovery rate	98%	> 75%	②
Default rate	0%	< 0%	8	Default rate	0%	< 15%	②

Death rate

Referral rate

Water, Sanitation and Hygiene (WASH)

0%

0%

< 10%

INDICATOR

100%

0.16

STANDARD

100%

> 0.5

PREVENTION & TREATMENT

Proportion of HIV positive mothers who

Number of PoCs receving ART

INDICATOR

60

STANDARD

100%

< 3%

2%

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	32	< 20	•	Number of hygiene promoters	10	1:<20,000	•
Proportion of households collecting >= 15 litres per person per day	Not	Not collected		SANITATION Refugees per latrine/toilet	INDICATOR 6	STANDARD < 20	②
Number of persons of concern per water tap	Not collected			Proportion of households with drop-hole latrine or drop-hole toilet	Not collected		
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5				Proportion of households report defecating in a toilet	Not collected		
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/	Not collected			Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	Not collected		
Proportion of households collecting drinking water from protected water sources only	Not	collected					
Proportion of households with sufficient daily water storage capacity	Not	collected					