

Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARD	
Number of medical doctors	0	1 : < 50,000	•	Incidence of malaria among children under 5	2		
Number of qualified nurses	5	1 : < 10,000	0	Incidence of watery diarrhoea among	19		
Number of community health workers	18	1 : < 1,000	▲	children under 5			
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	31		
Consultations per trained clinician per day	20	< 50	0	Incidence of pneumonia among over 5	10		
Health utilisation rate (new visits / refugee /	0.7	1 - 4	0	Tuberculosis success rate	100%	< 90%	
year) Proportion of host population consultations	24%			Were any MDR/X-TB cases diagnosed among PoCs?	No	No 😫	
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATO	RSTANDARD	
Crude Mortality Rate (CMR) (/1000/month)	0.1	< 0.75	0	Full vaccination coverage	92%	< 95%	▲
Under-five Mortality Rate (U5MR) (/1000/month)	0.1	< 1.5	0	Measles vaccination coverage	92%	< 95%	▲
Infant Mortality Rate (IMR) (/1000 livebirths)	1.7	< 30	0	Source of vaccination data	HIS		
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	0	IN-PATIENT DEPARTMENT (IPD)	INDICATO	R	
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	1.0		
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0		
Number of outbreaks reported	0			Hospitalisation rate	35.2	50 - 150	8

Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	61%	> 90%	8	Contraceptive prevalence rate	1%	≥ 30%	8
Coverage of antenatal tetanus vaccination	92%	> 95%	4	Source of contraceptive prevalence data	HIS		
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	90%	≥ 90%	4	Total number of reported rapes	1		
Proportion of births conducted by caesarean section	0%	5 - 15%	⊗	Proportion of eligible rape survivors provided with PEP within 72 hours	100%	100%	0
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	1%	< 15%	0	Proportion of eligible rape survivors provided with ECP within 120 hours	0%	100%	⊗
Proportion of teenage pregnancies	2%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate (/ 1000 total births / month)	0			Number of genital ulcer diseases	27		
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	41		
Proportion of maternal deaths investigated within 48 hours		100%					

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD	PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%	Number of PoCs receving ART	24	
Condom distribution rate	0.04	> 0.5	 Proportion of HIV positive mothers who swallowed ARV during delivery 		100%
Proportion of TB patients tested for HIV		100%	PMTCT coverage	96%	100%

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD		INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD	
Prevalence of global acute malnutrition (6-59 months)	Not collected			Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected		
Prevalence of stunting (6-59 months)	Not collected			Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected		
Prevalence of severe acute malnutrition (6 - 59 months)	Not collected			IYCF programmes targeting pregnant and lactating women established or maintained?	Yes		
Prevalence of anaemia in children (6-59 months)	Not collected			MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected			Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	Not collected		
CMAM: SAM TREATMENT	INDICATOR	STANDARD		MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)	102			Number of new moderate acute admissions (6-59 months)	520		
Recovery rate	98%	> 75%	0	Recovery rate	99%	> 75%	⊘
Default rate	1%	< 0%	8	Default rate	0%	< 15%	⊘
Death rate	1%	< 10%	0	Death rate	0%	< 3%	0
Referral rate	0%			Referral rate	1%		

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	14	< 20	8	Number of hygiene promoters		1:<20,000	•
Proportion of households collecting >= 15	Not collected			SANITATION	INDICATOR	STANDARD	
litres per person per day				Refugees per latrine/toilet	12	< 20	~
Number of persons of concern per water tap	Not collected			Proportion of households with drop-hole latrine or drop-hole toilet	Not		
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5	Not collected			· Proportion of households report defecating in a toilet	Not collected		
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	Not collected			Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	Not	collected	
Proportion of households collecting drinking water from protected water sources only	Not	collected					
Proportion of households with sufficient daily water storage capacity	Not	collected					