Site report 2018 Touloum CHAD

Population

24,282 refugees







Origin of refugees

SOUDAN

Site opened Jan 2004

Partners







Key Indicators

CRUDE MORTALITY RATE

0.1

Deaths/1,000 refugees/ month

GAM

Global Acute Malnutrition **U5 MORTALITY RATE**

0.2

Deaths/1,000 refugees/

SAM

NC

Severe Acute Malnutrition MATERNAL MORTALITY

0

Deaths

WATER SUPPLY

14

Liters/ person/ day

SKILLED BIRTH ATTENDANCE

97%


of all births


LATRINE COVERAGE

13


Refugees/communal toilets or latrines

COMMUNICABLE DISEASES





NON-COMMUNICABLE DISEASES



Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARD	
Number of medical doctors	0	1:<50,000	8	Incidence of malaria among children under 5	2		
Number of qualified nurses	4	1 : < 10,000	•	Incidence of watery diarrhoea among	17		
Number of community health workers	19	1:<1,000	8	children under 5			
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	46		
Consultations per trained clinician per day	20	< 50	•	Incidence of pneumonia among over 5	15		
Health utilisation rate (new visits / refugee /	0.7	1 - 4	8	Tuberculosis success rate	100%	< 90%	②
year) Proportion of host population consultations	7%			Were any MDR/X-TB cases diagnosed among PoCs?	No	No	8
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATOR	STANDA	RD
Crude Mortality Rate (CMR) (/1000/month)	0.1	< 0.75	②	Full vaccination coverage	139%	< 95%	•
Under-five Mortality Rate (U5MR) (/1000/month)	0.2	< 1.5	②	Measles vaccination coverage	139%	< 95%	,
Infant Mortality Rate (IMR) (/1000 livebirths)	7.6	< 30	②	Source of vaccination data	HIS		
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	②	IN-PATIENT DEPARTMENT (IPD)	INDICATOR	STANDA	RD
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	1.1		
Proportion of outbreaks investigated within 48 hours	100%	100%	②	Case fatality rate	0.0		
Number of outbreaks reported	2			Hospitalisation rate	35.7	50 - 15	0

Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	78%	> 90%	8	Contraceptive prevalence rate	15%	≥ 30%	(
Coverage of antenatal tetanus vaccination	94%	> 95%	<u> </u>	Source of contraceptive prevalence data	HIS		
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	97%	≥ 90%	②	Total number of reported rapes	0		
Proportion of births conducted by caesarean section	0%	5 - 15%	8	Proportion of eligible rape survivors provided with PEP within 72 hours		100%	
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	1%	< 15%	②	Proportion of eligible rape survivors provided with ECP within 120 hours		100%	
Proportion of teenage pregnancies	1%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate (/ 1000 total births / month)	1			Number of genital ulcer diseases	0		
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	0		
Proportion of maternal deaths investigated within 48 hours		100%					

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD	PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%	Number of PoCs receving ART	39	
Condom distribution rate	0.03	> 0.5	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%	PMTCT coverage	100%	100%

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD		INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD		
Prevalence of global acute malnutrition (6-59 months)	Not collected			Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected			
Prevalence of stunting (6-59 months)	Not collected			Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected			
Prevalence of severe acute malnutrition (6 - 59 months)	Not collected			IYCF programmes targeting pregnant and lactating women established or maintained?		Yes		
Prevalence of anaemia in children (6-59 months)	Not collected			MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD		
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected			Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	Not collected			
CMAM: SAM TREATMENT	INDICATOR	STANDARD		MAM TREATMENT	INDICATOR	STANDARD		
Number of new severe acute admissions (6-59 months)	251			Number of new moderate acute admissions (6-59 months)	859			
Recovery rate	97%	> 75%	②	Recovery rate	99%	> 75%	,	
Default rate	2%	< 0%	8	Default rate	0%	< 15%	,	
Death rate	0%	< 10%	②	Death rate	0%	< 3%	,	
Referral rate	1%			Referral rate	1%			

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD		
Average number of litres of potable water available per person per day	14	< 20	8	Number of hygiene promoters		1:<20,000	•	
	Not	collected		SANITATION	INDICATOR	STANDARD		
Proportion of households collecting >= 15 litres per person per day	Not collected			Refugees per latrine/toilet	13	< 20	②	
Number of persons of concern per water tap	Not collected			Proportion of households with drop-hole latrine or drop-hole toilet	Not collected			
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5	Not collected			Proportion of households report defecating in a toilet	Not collected			
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	Not collected			Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	Not	collected		
Proportion of households collecting drinking water from protected water sources only	Not	collected						
Proportion of households with sufficient daily water storage capacity	Not	collected						