Site report 2018 Ban Don Yang THAILAND

Population

2,449 refugees







Origin of refugees

MYANMAR

Site opened Jan 1997

Partners





Key Indicators

CRUDE MORTALITY RATE

Deaths/1,000 refugees/ month

GAM

Global Acute

U5 MORTALITY RATE

Deaths/1,000 refugees/ month

SAM

NC

Severe Acute

MATERNAL MORTALITY

Deaths

WATER SUPPLY

70

Liters/ person/ day


SKILLED BIRTH ATTENDANCE


102% of all births


LATRINE COVERAGE

Refugees/communal


COMMUNICABLE DISEASES







NON-COMMUNICABLE DISEASES



Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARD	
Number of medical doctors	1	1:<50,000	②	Incidence of malaria among children under 5	0		
Number of qualified nurses	1	1 : < 10,000	②	Incidence of watery diarrhoea among	36		
Number of community health workers	10	1 : < 1,000	②	children under 5			
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	173		
Consultations per trained clinician per day	21	< 50	②	Incidence of pneumonia among over 5	34		
Health utilisation rate (new visits / refugee /	2.7	1 - 4	②	Tuberculosis success rate	100%	< 90%	•
year) Proportion of host population consultations	15%			Were any MDR/X-TB cases diagnosed among PoCs?	No	No	8
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATOR	STANDA	ARD
Crude Mortality Rate (CMR) (/1000/month)	0.3	< 0.75	⊘	Full vaccination coverage	110%	< 95%	
Under-five Mortality Rate (U5MR) (/1000/month)	0.0	< 1.5	②	Measles vaccination coverage	106%	< 95%	6
Infant Mortality Rate (IMR) (/1000 livebirths)	0.0	< 30	②	Source of vaccination data	HIS		
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	②	IN-PATIENT DEPARTMENT (IPD)	INDICATOR	STANDA	ARD
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	4.8		
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0		
Number of outbreaks reported	0			Hospitalisation rate	153.9	50 - 15	50

Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	98%	> 90%	②	Contraceptive prevalence rate	48%	≥ 30%	1
Coverage of antenatal tetanus vaccination	100%	> 95%	②	Source of contraceptive prevalence data	HIS		
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	102%	≥ 90%	②	Total number of reported rapes	0		
Proportion of births conducted by caesarean section	11%	5 - 15%	②	Proportion of eligible rape survivors provided with PEP within 72 hours		100%	
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	2%	< 15%	②	Proportion of eligible rape survivors provided with ECP within 120 hours		100%	
Proportion of teenage pregnancies	11%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate (/ 1000 total births / month)	2			Number of genital ulcer diseases	0		
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	0		
Proportion of maternal deaths investigated within 48 hours		100%					

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD	PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%	Number of PoCs receving ART	2	
Condom distribution rate	0.20	> 0.5	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%	PMTCT coverage	100%	100%

Nutrition

NUTRITION GENERAL	INDICATOR STANDARD	INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD	
Prevalence of global acute malnutrition (6-59 months)	Not collected	Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected		
Prevalence of stunting (6-59 months)	Not collected	Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected		
Prevalence of severe acute malnutrition (6 - 59 months)	Not collected	IYCF programmes targeting pregnant and lactating women established or maintained?	Yes		
Prevalence of anaemia in children (6-59 months)	Not collected	MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected	Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	Not collected		
CMAM: SAM TREATMENT	INDICATOR STANDARD	MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)		Number of new moderate acute admissions (6-59 months)	5		
Recovery rate	> 75%	Recovery rate	80%	> 75%	
Default rate	< 0%	Default rate	0%	< 15%	
Death rate	< 10%	Death rate	0%	< 3%	
Referral rate		Referral rate	20%		

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD		
Average number of litres of potable water available per person per day	70	< 20	②	Number of hygiene promoters	10	1:<20,000	②	
Proportion of households collecting >= 15 litres per person per day	100%	> 95%	②	SANITATION Refugees per latrine/toilet	INDICATOR 5	STANDARD < 20	②	
Number of persons of concern per water tap	34	< 80	•	Proportion of households with drop-hole	100%	120		
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5	100%	> 95%	•	Proportion of households report defecating in a toilet		> 85%		
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%		Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%		
Proportion of households collecting drinking water from protected water sources only		> 95%						
Proportion of households with sufficient daily water storage capacity	100%	> 80%	②					