
Site report 2018 Ban Mae Surin THAILAND

Population

2,209 refugees







Origin of refugees

MYANMAR

Site opened Nov 1991

Partners





Key Indicators

CRUDE MORTALITY RATE

Deaths/1,000 refugees/ month

GAM

Global Acute

U5 MORTALITY RATE

Deaths/1,000 refugees/ month

SAM

NC

Severe Acute

MATERNAL MORTALITY

Deaths

WATER SUPPLY


15 Liters/ person/ day SKILLED BIRTH ATTENDANCE


100% of all births

LATRINE COVERAGE


Refugees/communal


COMMUNICABLE DISEASES





NON-COMMUNICABLE DISEASES





Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARD		
Number of medical doctors	2	1: < 50,000	②	Incidence of malaria among children under 5	0			
Number of qualified nurses	1	1:<10,000	②	Incidence of watery diarrhoea among	56			
Number of community health workers	7	1:<1,000	•	children under 5				
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	108			
Consultations per trained clinician per day	17	< 50	②	Incidence of pneumonia among over 5	19			
Health utilisation rate (new visits / refugee /	3.1	1 - 4	②	Tuberculosis success rate	100%	< 90%	•	
year)				Were any MDR/X-TB cases diagnosed among	No	No	8	
Proportion of host population consultations	0%			PoCs?				
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATO	R	ARD	
Crude Mortality Rate (CMR) (/1000/month)	0.3	< 0.75	⊘	Full vaccination coverage	91%	< 95	%	A
Under-five Mortality Rate (U5MR) (/1000/month)	0.6	< 1.5	②	Measles vaccination coverage	91%	< 95	%	A
Infant Mortality Rate (IMR) (/1000 livebirths)	46.5	< 30	8	Source of vaccination data	HIS			
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	47	< 20	8	IN-PATIENT DEPARTMENT (IPD)	INDICATO	R STAND	ARD	
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	4.3			
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0			
Number of outbreaks reported	0			Hospitalisation rate	50.7	50 - 1	50	②

Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	93%	> 90%	②	Contraceptive prevalence rate	48%	≥ 30%	(
Coverage of antenatal tetanus vaccination	98%	> 95%	②	Source of contraceptive prevalence data	HIS		
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	100%	≥ 90%	②	Total number of reported rapes	0		
Proportion of births conducted by caesarean section	9%	5 - 15%	②	Proportion of eligible rape survivors provided with PEP within 72 hours		100%	
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	9%	< 15%	②	Proportion of eligible rape survivors provided with ECP within 120 hours		100%	
Proportion of teenage pregnancies	0%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate (/ 1000 total births / month)	0			Number of genital ulcer diseases	0		
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	0		
Proportion of maternal deaths investigated within 48 hours		100%					

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD		PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%		Number of PoCs receving ART	1	
Condom distribution rate	0.01	> 0.5	8	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%		PMTCT coverage	100%	100%

Nutrition

NUTRITION GENERAL	INDICATOR STANDARD	INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD	
NOTATION GENERAL	INDICATOR	INFANT AND TOUNG CHIED FEEDING (ITCF)	INDICATOR	STANDARD	
Prevalence of global acute malnutrition (6-59 months)	Not collected	Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected		
Prevalence of stunting (6-59 months)	Not collected	Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected		
Prevalence of severe acute malnutrition (6 - 59 months)	Not collected	IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	
Prevalence of anaemia in children (6-59 months)	Not collected	MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected	Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	Not collected		
CMAM: SAM TREATMENT	INDICATOR STANDARD	MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)		Number of new moderate acute admissions (6-59 months)	6		
Recovery rate	> 75%	Recovery rate	33%	> 75%	8
Default rate	< 0%	Default rate	7%	< 15%	②
Death rate	< 10%	Death rate	0%	< 3%	②
Referral rate		Referral rate	0%		

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	15	< 20	8	Number of hygiene promoters		1:<20,000	•
Proportion of households collecting >= 15 litres per person per day		> 95%		SANITATION Defrages per letring /tailet	INDICATOR 5	STANDARD < 20	•
Number of persons of concern per water tap	5	< 80	•	Refugees per latrine/toilet Proportion of households with drop-hole	5	< 20	
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU		> 95%		latrine or drop-hole toilet Proportion of households report defecating in a toilet		> 85%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%		Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%	
Proportion of households collecting drinking water from protected water sources only		> 95%					
Proportion of households with sufficient daily water storage capacity		> 80%					