
Site report 2018 Mae La Oon THAILAND

Population

10,206 refugees







Origin of refugees

MYANMAR

Site opened Jan 2004

Partners

Key Indicators

CRUDE MORTALITY RATE

Deaths/1,000 refugees/

GAM

Global Acute Malnutrition **U5 MORTALITY RATE**

Deaths/1,000 refugees/

SAM

NC

Severe Acute Malnutrition

MATERNAL MORTALITY

Deaths

WATER SUPPLY

Liters/ person/ day

SKILLED BIRTH **ATTENDANCE**


78% of all births


LATRINE COVERAGE

Refugees/communal toilets or latrines


COMMUNICABLE DISEASES

15.8 %





NON-COMMUNICABLE DISEASES



Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARD	
Number of medical doctors		1: < 50,000	•	Incidence of malaria among children under 5	1		
Number of qualified nurses		1:<10,000	•	Incidence of watery diarrhoea among	57		
Number of community health workers		1:<1,000	•	children under 5			
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	106		
Consultations per trained clinician per day	30	< 50	②	Incidence of pneumonia among over 5	25		
Health utilisation rate (new visits / refugee /	2.1	1 - 4	•	Tuberculosis success rate		< 90%	
year) Proportion of host population consultations	5%			Were any MDR/X-TB cases diagnosed among PoCs?		No	
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATO	STANDAR	D
Crude Mortality Rate (CMR) (/1000/month)	0.4	< 0.75	②	Full vaccination coverage	62%	< 95%	8
Under-five Mortality Rate (U5MR) (/1000/month)	0.3	< 1.5	②	Measles vaccination coverage	62%	< 95%	8
Infant Mortality Rate (IMR) (/1000 livebirths)	12.5	< 30	②	Source of vaccination data			
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	②	IN-PATIENT DEPARTMENT (IPD)	INDICATO	STANDAR	D
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	5.0		
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0		
Number of outbreaks reported	0			Hospitalisation rate	153.5	50 - 150	O

Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	94%	> 90%	②	Contraceptive prevalence rate	0%	≥ 30%	
Coverage of antenatal tetanus vaccination	98%	> 95%	②	Source of contraceptive prevalence data			
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	78%	≥ 90%	8	Total number of reported rapes	0		
Proportion of births conducted by caesarean section	4%	5 - 15%	A	Proportion of eligible rape survivors provided with PEP within 72 hours		100%	
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	8%	< 15%	②	Proportion of eligible rape survivors provided with ECP within 120 hours		100%	
Proportion of teenage pregnancies	1%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate (/ 1000 total births / month)	1			Number of genital ulcer diseases	0		
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	0		
Proportion of maternal deaths investigated within 48 hours		100%					

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD		PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV	100%	100%	②	Number of PoCs receving ART	1	
Condom distribution rate	0.00	> 0.5	8	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%		PMTCT coverage	100%	100%

Nutrition

NUTRITION GENERAL	INDICATOR STANDARD	INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD	
Prevalence of global acute malnutrition (6-59 months)	Not collected	Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected		
Prevalence of stunting (6-59 months)	Not collected	Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected		
Prevalence of severe acute malnutrition (6 - 59 months)	Not collected	IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	
Prevalence of anaemia in children (6-59 months)	Not collected	MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected	Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)		> 70%	
CMAM: SAM TREATMENT	INDICATOR STANDARD	MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)		Number of new moderate acute admissions (6-59 months)	20		
Recovery rate	> 75%	Recovery rate	85%	> 75%	
Default rate	< 0%	Default rate	0%	< 15%	②
Death rate	< 10%	Death rate	0%	< 3%	②
Referral rate		Referral rate	10%		

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD	WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day		< 20	Number of hygiene promoters		1:<20,000	•
Proportion of households collecting >= 15		> 95%	SANITATION	INDICATOR	STANDARD	
litres per person per day			Refugees per latrine/toilet		< 20	
Number of persons of concern per water tap		< 80	Proportion of households with drop-hole latrine or drop-hole toilet			
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5		> 95%	Proportion of households report defecating in a toilet		> 85%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%	Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%	
Proportion of households collecting drinking water from protected water sources only		> 95%				
Proportion of households with sufficient daily water storage capacity		> 80%				