# Site report 2018 Mae Ra Ma Luang THAILAND






**Origin of refugees MYANMAR** 


Site opened Feb 1995


#### Partners

## **Key Indicators**

#### SKILLED BIRTH CRUDE MORTALITY RATE **U5 MORTALITY RATE** MATERNAL MORTALITY ATTENDANCE Deaths/1,000 refugees/ Deaths/1,000 refugees/ 0.30.3 79% of all births 0 Deaths month month GAM SAM WATER SUPPLY LATRINE COVERAGE Severe Acute Global Acute NC Refugees/communal NC Liters/ person/ day Malnutrition Malnutrition toilets or latrines COMMUNICABLE DISEASES **NON-COMMUNICABLE DISEASES** 47.0 % 13.8 % LRTI 23.4 % URTI 13.0 % Skin disease 13.8 % LRTI 23.4 % 6.8 % URTI 20.4 % 13.0 % 21.0 % Respiratory disorders Skin Other Total disease 43.0 % 11.6 % Consultations Other 25,584 6.8 % **MENTAL HEALTH** INJURIES 66.3 %







## Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARD	
Number of medical doctors		1 : < 50,000	•	Incidence of malaria among children under 5	0		
Number of qualified nurses		1 : < 10,000	•	Incidence of watery diarrhoea among	31		
Number of community health workers		1 : < 1,000	•	children under 5			
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	120		
Consultations per trained clinician per day	31	< 50	0	Incidence of pneumonia among over 5	29		
Health utilisation rate (new visits / refugee /	2.5	1 - 4	0	Tuberculosis success rate		< 90%	
year)				Were any MDR/X-TB cases diagnosed among PoCs?		No	
Proportion of host population consultations	2%			POCS?			
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATO	STANDARD	
Crude Mortality Rate (CMR) (/1000/month)	0.3	< 0.75	0	Full vaccination coverage	81%	< 95%	8
Under-five Mortality Rate (U5MR) (/1000/month)	0.3	< 1.5	<b>~</b>	Measles vaccination coverage	82%	< 95%	⊗
Infant Mortality Rate (IMR) (/1000 livebirths)	12.6	< 30	0	Source of vaccination data			
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	8	< 20	0	IN-PATIENT DEPARTMENT (IPD)	INDICATO	STANDARD	
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	5.0		
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0		
Number of outbreaks reported	0			Hospitalisation rate	171.0	50 - 150	0

# Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	94%	> 90%	0	Contraceptive prevalence rate	0%	≥ 30%	8
Coverage of antenatal tetanus vaccination	97%	> 95%	<b>~</b>	Source of contraceptive prevalence data			
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	79%	≥ 90%	⊗	Total number of reported rapes	1		
Proportion of births conducted by caesarean section	5%	5 - 15%	0	Proportion of eligible rape survivors provided with PEP within 72 hours		100%	
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	6%	< 15%	0	Proportion of eligible rape survivors provided with ECP within 120 hours		100%	
Proportion of teenage pregnancies	2%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate ( / 1000 total births / month)	2			Number of genital ulcer diseases	0		
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	0		
Proportion of maternal deaths investigated within 48 hours		100%					

### HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD		PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV	100%	100%	0	Number of PoCs receving ART	0	
Condom distribution rate	0.00	> 0.5	8	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%		PMTCT coverage	97%	100%

# Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD	INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD	
Prevalence of global acute malnutrition (6-59 months)	Not collected		Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected		
Prevalence of stunting (6-59 months)	Not collected		Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected		
Prevalence of severe acute malnutrition (6 - 59 months)	Not collected		IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	
Prevalence of anaemia in children (6-59 months)	Not collected		MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected		Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)		> 70%	
CMAM: SAM TREATMENT	INDICATOR	STANDARD	MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)			Number of new moderate acute admissions (6-59 months)	28		
Recovery rate		> 75%	Recovery rate	85%	> 75%	<b>S</b>
Default rate		< 0%	Default rate	0%	< 15%	<b>S</b>
Death rate		< 10%	Death rate	0%	< 3%	0
Referral rate			Referral rate	15%		

# Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD	WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day		< 20	Number of hygiene promoters		1 : < 20,000	•
Proportion of households collecting >= 15		> 95%	SANITATION	INDICATOR	STANDARD	
litres per person per day			Refugees per latrine/toilet		< 20	
Number of persons of concern per water tap		< 80	Proportion of households with drop-hole latrine or drop-hole toilet			
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual ( >= 0.1 mg/litre) and NTU <5		> 95%	Proportion of households report defecating in a toilet		> 85%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%	Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%	
Proportion of households collecting drinking water from protected water sources only		> 95%				
Proportion of households with sufficient daily water storage capacity		> 80%				