
# Site report 2018 Nu Poh THAILAND

## **Population**

9,042 refugees







#### Origin of refugees

**MYANMAR** 

Site opened Mar 1997

#### **Partners**





## **Key Indicators**

**CRUDE MORTALITY RATE** 

0.3

GAM

Deaths/1,000 refugees/ month

Global Acute

U5 MORTALITY RATE

0.2

Deaths/1,000 refugees/

SAM

NC

Severe Acute Malnutrition MATERNAL MORTALITY

n

Deaths

WATER SUPPLY

Liters/ person/ day

SKILLED BIRTH ATTENDANCE

99%


of all births


LATRINE COVERAGE

51.7 %


Refugees/communal toilets or latrines

#### **COMMUNICABLE DISEASES**





#### **NON-COMMUNICABLE DISEASES**



## **Public Health**

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARD	
Number of medical doctors	1	1:<50,000	•	Incidence of malaria among children under 5	0		
Number of qualified nurses	21	1:<10,000	<b>②</b>	Incidence of watery diarrhoea among	43		
Number of community health workers	23	1:<1,000	•	children under 5			
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	129		
Consultations per trained clinician per day	20	< 50	<b>②</b>	Incidence of pneumonia among over 5	31		
Health utilisation rate (new visits / refugee /	1.9	1 - 4	•	Tuberculosis success rate	85%	< 90%	<b>A</b>
year)  Proportion of host population consultations	18%			Were any MDR/X-TB cases diagnosed among PoCs?	No	No	8
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATOR	STANDAR	RD
Crude Mortality Rate (CMR) (/1000/month)	0.3	< 0.75	•	Full vaccination coverage	105%	< 95%	<b>②</b>
Under-five Mortality Rate (U5MR) (/1000/month)	0.2	< 1.5	<b>②</b>	Measles vaccination coverage	117%	< 95%	<b>⊘</b>
Infant Mortality Rate (IMR) (/1000 livebirths)	0.0	< 30	<b>②</b>	Source of vaccination data	HIS		
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	<b>②</b>	IN-PATIENT DEPARTMENT (IPD)	INDICATOR	STANDAR	RD
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	7.0		
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0		
Number of outbreaks reported	0			Hospitalisation rate	137.2	50 - 150	<b>O</b>

## **Reproductive Health**

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	93%	> 90%	<b>②</b>	Contraceptive prevalence rate	49%	≥ 30%	
Coverage of antenatal tetanus vaccination	96%	> 95%	<b>②</b>	Source of contraceptive prevalence data	HIS		
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	99%	≥ 90%	<b>⊘</b>	Total number of reported rapes	0		
Proportion of births conducted by caesarean section	16%	5 - 15%	<b>②</b>	Proportion of eligible rape survivors provided with PEP within 72 hours		100%	
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	9%	< 15%	<b>②</b>	Proportion of eligible rape survivors provided with ECP within 120 hours		100%	
Proportion of teenage pregnancies	5%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate ( / 1000 total births / month)	0			Number of genital ulcer diseases	0		
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	11		
Proportion of maternal deaths investigated within 48 hours		100%					

### **HIV/AIDS**

PREVENTION & TREATMENT	INDICATOR	STANDARD		PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV	100%	100%	<b>②</b>	Number of PoCs receving ART	13	
Condom distribution rate	0.10	> 0.5	8	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%		PMTCT coverage	99%	100%

## Nutrition

NUTRITION GENERAL	INDICATOR STANDARD	INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD	
Prevalence of global acute malnutrition (6-59 months)	Not collected	Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected		
Prevalence of stunting (6-59 months)	Not collected	Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected		
Prevalence of severe acute malnutrition (6 - 59 months)	Not collected	IYCF programmes targeting pregnant and lactating women established or maintained?	Yes		
Prevalence of anaemia in children (6-59 months)	Not collected	MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected	Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	Not collected		
CMAM: SAM TREATMENT	INDICATOR STANDARD	MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)		Number of new moderate acute admissions (6-59 months)	4		
Recovery rate	> 75%	Recovery rate	18%	> 75%	3
Default rate	< 0%	Default rate	0%	< 15%	3
Death rate	< 10%	Death rate	0%	< 3%	<b>3</b>
Referral rate		Referral rate	0%		

## Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day		< 20		Number of hygiene promoters	35	1:<20,000	•
Proportion of households collecting >= 15 litres per person per day	100%	> 95%	•	SANITATION  Refugees per latrine/toilet	INDICATOR	STANDARD < 20	
Number of persons of concern per water tap	28	< 80	•	Proportion of households with drop-hole latrine or drop-hole toilet	100%		
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual ( >= 0.1 mg/litre) and NTU <5	100%	> 95%	<b>⊘</b>	Proportion of households report defecating in a toilet		> 85%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	100%	> 95%	•	Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	100%	> 90%	•
Proportion of households collecting drinking water from protected water sources only		> 95%					
Proportion of households with sufficient daily water storage capacity		> 80%					