# Site report 2018 Tham Hin THAILAND

### **Population**

5,238 refugees







#### **Origin of refugees**

**MYANMAR** 

Site opened May 1997

#### **Partners**





#### **Key Indicators**

**CRUDE MORTALITY RATE** 

Deaths/1,000 refugees/ month

**GAM** 

Global Acute

6.5 %

11.5 %

**U5 MORTALITY RATE** 

Deaths/1,000 refugees/ month

SAM

NC Severe Acute MATERNAL MORTALITY

Deaths

WATER SUPPLY

54

Liters/ person/ day


SKILLED BIRTH ATTENDANCE


100% of all births

LATRINE COVERAGE


Refugees/communal toilets or latrines


#### **COMMUNICABLE DISEASES**





#### **NON-COMMUNICABLE DISEASES**





## Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARD		
Number of medical doctors	1	1:<50,000	<b>Ø</b>	Incidence of malaria among children under 5	0			
Number of qualified nurses	1	1 : < 10,000	<b>②</b>	Incidence of watery diarrhoea among	47			
Number of community health workers	20	1:<1,000	•	children under 5				
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	232			
Consultations per trained clinician per day	31	< 50	•	Incidence of pneumonia among over 5	38			
Health utilisation rate (new visits / refugee /	2.9	1 - 4	•	Tuberculosis success rate	100%	< 90%	•	
year)  Proportion of host population consultations	0%			Were any MDR/X-TB cases diagnosed among PoCs?	No	No	8	
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATOR	STAND	ARD	
Crude Mortality Rate (CMR) (/1000/month)	0.3	< 0.75	<b>②</b>	Full vaccination coverage	118%	< 95	%	•
Under-five Mortality Rate (U5MR) (/1000/month)	0.0	< 1.5	•	Measles vaccination coverage	121%	< 95	%	•
Infant Mortality Rate (IMR) (/1000 livebirths)	0.0	< 30	<b>3</b>	Source of vaccination data	HIS			
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	<b>3</b>	IN-PATIENT DEPARTMENT (IPD)	INDICATOR	STAND	ARD	
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	6.5			
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0			
Number of outbreaks reported	0			Hospitalisation rate	137.5	50 - 1	50	•

# Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	98%	> 90%	<b>②</b>	Contraceptive prevalence rate	48%	≥ 30%	(
Coverage of antenatal tetanus vaccination	99%	> 95%	<b>②</b>	Source of contraceptive prevalence data	HIS		
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	100%	≥ 90%	<b>②</b>	Total number of reported rapes	0		
Proportion of births conducted by caesarean section	16%	5 - 15%	<b>②</b>	Proportion of eligible rape survivors provided with PEP within 72 hours		100%	
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	7%	< 15%	<b>②</b>	Proportion of eligible rape survivors provided with ECP within 120 hours		100%	
Proportion of teenage pregnancies	12%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate ( / 1000 total births / month)	1			Number of genital ulcer diseases	1		
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	6		
Proportion of maternal deaths investigated within 48 hours		100%					

#### **HIV/AIDS**

1114771123						
PREVENTION & TREATMENT	INDICATOR	STANDARD		PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%		Number of PoCs receving ART	8	
Condom distribution rate	0.11	> 0.5	8	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%		PMTCT coverage	99%	100%
Nutrition						
NUTRITION GENERAL	INDICATOR	STANDARD		INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	Not collected			Proportion of children 0-23 months who were put to the breast within one hour of birth	Not	collected
Prevalence of stunting (6-59 months)	Not	collected		Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not	collected
Prevalence of severe acute malnutrition (6 - 59 months)	Not	collected		IYCF programmes targeting pregnant and lactating women established or maintained?		Yes
Prevalence of anaemia in children (6-59 months)	Not	collected		MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not	collected		Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	Not	collected

# Number of new severe acute admissions (6-59 months) Number of new moderate acute admissions (6-59 months) Recovery rate > 75% Recovery rate > 0 Default rate > 0 Default rate

MAM TREATMENT

Death rate < 10% Death rate < 3%

Referral rate Referral rate

**INDICATOR** 

## Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	54	< 20	•	Number of hygiene promoters	20	1:<20,000	•
Proportion of households collecting >= 15 litres per person per day	100%	> 95%	•	SANITATION  Refugees per latrine/toilet	INDICATOR 5	STANDARD < 20	<b>•</b>
Number of persons of concern per water tap	18	< 80	•	Proportion of households with drop-hole latrine or drop-hole toilet	100%		
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual ( >= 0.1 mg/litre) and NTU <5	100%	> 95%	<b>Ø</b>	Proportion of households report defecating in a toilet		> 85%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	0%	> 95%	8	Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)		> 90%	
Proportion of households collecting drinking water from protected water sources only		> 95%					
Proportion of households with sufficient daily water storage capacity	100%	> 80%	<b>②</b>				