
Site report 2018 Umpiem Mai THAILAND

Population

10,100 refugees







Origin of refugees

MYANMAR

Site opened Jan 1999

Partners





Key Indicators

CRUDE MORTALITY RATE

Deaths/1,000 refugees/ month

GAM

Global Acute

U5 MORTALITY RATE

Deaths/1,000 refugees/ month

SAM

NC Severe Acute MATERNAL MORTALITY

Deaths

WATER SUPPLY

Liters/ person/ day

SKILLED BIRTH ATTENDANCE


100% of all births


LATRINE COVERAGE

51.6 %


Refugees/communal toilets or latrines

COMMUNICABLE DISEASES





NON-COMMUNICABLE DISEASES



Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARD	
Number of medical doctors	1	1:<50,000	•	Incidence of malaria among children under	0		
Number of qualified nurses	26	1 : < 10,000	•	Incidence of watery diarrhoea among children under 5	16		
Number of community health workers ACCESS AND UTILIZATION	22 INDICATOR	1 : < 1,000 STANDARD	②	Incidence of pneumonia among children under 5	61		
Consultations per trained clinician per day	18	< 50	•	Incidence of pneumonia among over 5	10		
Health utilisation rate (new visits / refugee /	1.8	1 - 4	Ø	Tuberculosis success rate	100%	< 90%	
year) Proportion of host population consultations	1%			Were any MDR/X-TB cases diagnosed among PoCs?	No	No 6	9
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATO	R STANDARI	
Crude Mortality Rate (CMR) (/1000/month)	0.3	< 0.75	•	Full vaccination coverage	90%	< 95%	A
Under-five Mortality Rate (U5MR) (/1000/month)	0.0	< 1.5	②	Measles vaccination coverage	91%	< 95%	Δ
Infant Mortality Rate (IMR) (/1000 livebirths)	0.0	< 30	3	Source of vaccination data	HIS		
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	3	IN-PATIENT DEPARTMENT (IPD)	INDICATO	STANDARI	D .
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	7.4		
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0		
Number of outbreaks reported	0			Hospitalisation rate	133.8	50 - 150	②

Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	94%	> 90%	②	Contraceptive prevalence rate	56%	≥ 30%	•
Coverage of antenatal tetanus vaccination	97%	> 95%	②	Source of contraceptive prevalence data	HIS		
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	100%	≥ 90%	•	Total number of reported rapes	2		
Proportion of births conducted by caesarean section	17%	5 - 15%	②	Proportion of eligible rape survivors provided with PEP within 72 hours		100%	
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	6%	< 15%	②	Proportion of eligible rape survivors provided with ECP within 120 hours	100%	100%	•
Proportion of teenage pregnancies	5%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate (/ 1000 total births / month)	3			Number of genital ulcer diseases	1		
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	16		
Proportion of maternal deaths investigated within 48 hours		100%					

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD		PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV	100%	100%	②	Number of PoCs receving ART	18	
Condom distribution rate	0.10	> 0.5	8	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%		PMTCT coverage	100%	100%

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD	INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD		
Prevalence of global acute malnutrition (6-59 months)	Not collected		Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected			
Prevalence of stunting (6-59 months)	Not collected		Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected			
Prevalence of severe acute malnutrition (6 - 59 months)	Not collected		IYCF programmes targeting pregnant and lactating women established or maintained?		Yes		
Prevalence of anaemia in children (6-59 months)	Not collected		MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD		
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected		Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	Not collected			
CMAM: SAM TREATMENT	INDICATOR	STANDARD	MAM TREATMENT	INDICATOR	STANDARD		
Number of new severe acute admissions (6-59 months)			Number of new moderate acute admissions (6-59 months)	12			
Recovery rate		> 75%	Recovery rate	69%	> 75%	A	
Default rate		< 0%	Default rate	15%	< 15%	<u> </u>	
Death rate		< 10%	Death rate	0%	< 3%	Ø	
Referral rate			Referral rate	0%			

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD		
Average number of litres of potable water available per person per day		< 20		Number of hygiene promoters	47	1:<20,000	Ø	
Proportion of households collecting >= 15 litres per person per day	100%	> 95%	•	SANITATION Refugees per latrine/toilet	INDICATOR	STANDARD < 20		
Number of persons of concern per water tap	51	< 80	•	Proportion of households with drop-hole latrine or drop-hole toilet	60%			
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5	97%	> 95%	•	Proportion of households report defecating in a toilet		> 85%		
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	88%	> 95%	A	Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	100%	> 90%	•	
Proportion of households collecting drinking water from protected water sources only		> 95%						
Proportion of households with sufficient daily water storage capacity		> 80%						