

Site report 2018 Mtendeli TANZANIA



Population


41,640 refugees

20%

Children under 5

20%

Women of reproductive age



Origin of refugees

BURUNDI

Site opened
Jan 2016

Partners



Key Indicators

CRUDE MORTALITY RATE

0.1

Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.3

Deaths/1,000 refugees/month

MATERNAL MORTALITY

0

Deaths

SKILLED BIRTH ATTENDANCE

97%

of all births

GAM

3%

Global Acute Malnutrition

SAM

0

Severe Acute Malnutrition

WATER SUPPLY

20


Liters/ person/ day

LATRINE COVERAGE

8


Refugees/communal toilets or latrines

COMMUNICABLE DISEASES




INJURIES

NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING

	INDICATOR	STANDARD
Number of medical doctors	4	1 : < 50,000
Number of qualified nurses	27	1 : < 10,000
Number of community health workers	1	1 : < 1,000

ACCESS AND UTILIZATION

	INDICATOR	STANDARD
Consultations per trained clinician per day	30	< 50
Health utilisation rate (new visits / refugee / year)	1.4	1 - 4
Proportion of host population consultations	0%	

MORTALITY

	INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.1	< 0.75
Under-five Mortality Rate (U5MR) (/1000/month)	0.3	< 1.5
Infant Mortality Rate (IMR) (/1000 livebirths)	8.9	< 30
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	3	< 20

OUTBREAK ALERT AND RESPONSE

	INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours		100%
Number of outbreaks reported	0	

MORBIDITY

	INDICATOR	STANDARD
Incidence of malaria among children under 5	72	
Incidence of watery diarrhoea among children under 5	4	
Incidence of pneumonia among children under 5	14	
Incidence of pneumonia among over 5	9	
Tuberculosis success rate	75%	< 90%
Were any MDR/X-TB cases diagnosed among PoCs?	No	No

VACCINATION

	INDICATOR	STANDARD
Full vaccination coverage	80%	< 95%
Measles vaccination coverage	99%	< 95%
Source of vaccination data	HIS	

IN-PATIENT DEPARTMENT (IPD)

	INDICATOR	STANDARD
Average length of stay (days)	1.8	
Case fatality rate	0.0	
Hospitalisation rate	101.3	50 - 150

Reproductive Health

ANTENATAL CARE

	INDICATOR	STANDARD
Antenatal care coverage	86%	> 90%
Coverage of antenatal tetanus vaccination	93%	> 95%

DELIVERY CARE

	INDICATOR	STANDARD
Proportion of births attended by skilled personnel	97%	≥ 90%
Proportion of births conducted by caesarean section	12%	5 - 15%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	9%	< 15%
Proportion of teenage pregnancies	2%	
Still birth rate (/ 1000 total births / month)	1	
Number of maternal deaths	0	
Proportion of maternal deaths investigated within 48 hours		100%

FAMILY PLANNING

	INDICATOR	STANDARD
Contraceptive prevalence rate	30%	≥ 30%
Source of contraceptive prevalence data	HIS	

SEXUAL AND GENDER-BASED VIOLENCE (SGBV)

	INDICATOR	STANDARD
Total number of reported rapes	41	
Proportion of eligible rape survivors provided with PEP within 72 hours	90%	100%
Proportion of eligible rape survivors provided with ECP within 120 hours	62%	100%

SEXUALLY TRANSMITTED INFECTIONS (STIS)

	INDICATOR	STANDARD
Number of genital ulcer diseases	99	
Number of Pelvic Inflammatory diseases	5	

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD	PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV	100%	100%	✓ Number of PoCs receiving ART	50	
Condom distribution rate	0.01	> 0.5	✗ Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%	PMTCT coverage	106%	100% ✓

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD	INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	3%	< 20%	✓ Proportion of children 0-23 months who were put to the breast within one hour of birth	89%	> 80% ✓
Prevalence of stunting (6-59 months)	49%	< 20%	✓ Proportion of infants aged between 0-5 months who received only breast milk during the previous day	87%	> 70% ✓
Prevalence of severe acute malnutrition (6 - 59 months)	0%	< 3%	✓ IYCF programmes targeting pregnant and lactating women established or maintained?		Yes
Prevalence of anaemia in children (6-59 months)	50%	< 20%	✗ MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD
Prevalence of anaemia in women of reproductive age (15-49 yrs)	21%	< 20%	⚠ Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	90%	> 70% ✓
CMAM: SAM TREATMENT	INDICATOR	STANDARD	MAM TREATMENT	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	199		Number of new moderate acute admissions (6-59 months)	0	
Recovery rate	98%	> 75%	✓ Recovery rate		> 75%
Default rate	1%	< 0%	✗ Default rate		< 15%
Death rate	1%	< 10%	✓ Death rate		< 3%
Referral rate	0%		Referral rate		

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD	WATER STAFFING	INDICATOR	STANDARD
Average number of litres of potable water available per person per day	20	< 20	✓ Number of hygiene promoters	75	1 : < 20,000 ✓
Proportion of households collecting >= 15 litres per person per day	13%	> 95%	✗	INDICATOR	STANDARD
Number of persons of concern per water tap	92	< 80	⚠	Refugees per latrine/toilet	8 < 20 ✓
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5	100%	> 95%	✓	Proportion of households with drop-hole latrine or drop-hole toilet	41%
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	0%	> 95%	✗	Proportion of households report defecating in a toilet	100% > 85% ✓
Proportion of households collecting drinking water from protected water sources only	99%	> 95%	✓	Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	2% > 90% ✗
Proportion of households with sufficient daily water storage capacity	20%	> 80%	✗		