

# Site report 2018 Nduta TANZANIA



## Population


**96,634** refugees

**22%**

Children under 5

**20%**

Women of reproductive age



Origin of refugees

**BURUNDI**

Site opened  
Oct 2015

## Partners



## Key Indicators

CRUDE MORTALITY RATE

**0.1** Deaths/1,000 refugees/month

U5 MORTALITY RATE

**0.4** Deaths/1,000 refugees/month

MATERNAL MORTALITY

**0** Deaths

SKILLED BIRTH ATTENDANCE

**81%** of all births

GAM

**2%** Global Acute Malnutrition

SAM

**0** Severe Acute Malnutrition


WATER SUPPLY

**19** Liters/ person/ day


LATRINE COVERAGE

**5** Refugees/communal toilets or latrines


### COMMUNICABLE DISEASES




### INJURIES



### NON-COMMUNICABLE DISEASES



### MENTAL HEALTH



## Public Health

### HEALTH STAFFING

	INDICATOR	STANDARD
Number of medical doctors	9	1 : < 50,000
Number of qualified nurses	90	1 : < 10,000
Number of community health workers	136	1 : < 1,000

### ACCESS AND UTILIZATION

	INDICATOR	STANDARD
Consultations per trained clinician per day	42	< 50
Health utilisation rate (new visits / refugee / year)	2.1	1 - 4
Proportion of host population consultations	0%	

### MORTALITY

	INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.1	< 0.75
Under-five Mortality Rate (U5MR) (/1000/month)	0.4	< 1.5
Infant Mortality Rate (IMR) (/1000 livebirths)	19.0	< 30
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	15	< 20

### OUTBREAK ALERT AND RESPONSE

	INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours		100%
Number of outbreaks reported	0	

### MORBIDITY

	INDICATOR	STANDARD
Incidence of malaria among children under 5	45	
Incidence of watery diarrhoea among children under 5	54	
Incidence of pneumonia among children under 5	23	
Incidence of pneumonia among over 5	8	
Tuberculosis success rate	87%	< 90%
Were any MDR/X-TB cases diagnosed among PoCs?	No	No

### VACCINATION

	INDICATOR	STANDARD
Full vaccination coverage	0%	< 95%
Measles vaccination coverage	100%	< 95%
Source of vaccination data	HIS	

### IN-PATIENT DEPARTMENT (IPD)

	INDICATOR	STANDARD
Average length of stay (days)	3.9	
Case fatality rate	0.0	
Hospitalisation rate	104.8	50 - 150

## Reproductive Health

### ANTENATAL CARE

	INDICATOR	STANDARD
Antenatal care coverage	71%	> 90%
Coverage of antenatal tetanus vaccination	92%	> 95%

### DELIVERY CARE

	INDICATOR	STANDARD
Proportion of births attended by skilled personnel	81%	≥ 90%
Proportion of births conducted by caesarean section	2%	5 - 15%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	10%	< 15%
Proportion of teenage pregnancies	3%	
Still birth rate ( / 1000 total births / month)	1	
Number of maternal deaths	0	
Proportion of maternal deaths investigated within 48 hours		100%

### FAMILY PLANNING

	INDICATOR	STANDARD
Contraceptive prevalence rate	4%	≥ 30%
Source of contraceptive prevalence data	HIS	

### SEXUAL AND GENDER-BASED VIOLENCE (SGBV)

	INDICATOR	STANDARD
Total number of reported rapes	226	
Proportion of eligible rape survivors provided with PEP within 72 hours	88%	100%
Proportion of eligible rape survivors provided with ECP within 120 hours	55%	100%

### SEXUALLY TRANSMITTED INFECTIONS (STIS)

	INDICATOR	STANDARD
Number of genital ulcer diseases	0	
Number of Pelvic Inflammatory diseases	0	

## HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD	PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV	8%	100%	✓ Number of PoCs receiving ART	619	
Condom distribution rate	0.19	> 0.5	✗ Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%	PMTCT coverage	129%	100% ✓

## Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD	INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	2%	< 20%	✓ Proportion of children 0-23 months who were put to the breast within one hour of birth	88%	> 80% ✓
Prevalence of stunting (6-59 months)	57%	< 20%	⚠ Proportion of infants aged between 0-5 months who received only breast milk during the previous day	77%	> 70% ✓
Prevalence of severe acute malnutrition (6 - 59 months)	0%	< 3%	✓ IYCF programmes targeting pregnant and lactating women established or maintained?		Yes
Prevalence of anaemia in children (6-59 months)	37%	< 20%	✗ MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD
Prevalence of anaemia in women of reproductive age (15-49 yrs)	12%	< 20%	✓ Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/BBF)	90%	> 70% ✓
CMAM: SAM TREATMENT	INDICATOR	STANDARD	MAM TREATMENT	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	345		Number of new moderate acute admissions (6-59 months)	1566	
Recovery rate	92%	> 75%	✓ Recovery rate	91%	> 75% ✓
Default rate	5%	< 0%	✗ Default rate	1%	< 15% ✓
Death rate	1%	< 10%	✓ Death rate	0%	< 3% ✓
Referral rate	0%		Referral rate	2%	

## Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD	WATER STAFFING	INDICATOR	STANDARD
Average number of litres of potable water available per person per day	19	< 20	⚠ Number of hygiene promoters	106	1 : < 20,000 ✓
Proportion of households collecting >= 15 litres per person per day	11%	> 95%	✗	INDICATOR	STANDARD
Number of persons of concern per water tap	89	< 80	⚠ Refugees per latrine/toilet	5	< 20 ✓
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual ( >= 0.1 mg/litre) and NTU <5	100%	> 95%	✓ Proportion of households with drop-hole latrine or drop-hole toilet	44%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml		> 95%	✓ Proportion of households report defecating in a toilet	99%	> 85% ✓
Proportion of households collecting drinking water from protected water sources only	98%	> 95%	✓ Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	0%	> 90% ✗
Proportion of households with sufficient daily water storage capacity	55%	> 80%	✗		