



Site report 2018 Nyarugusu TANZANIA



Population

148,084 refugees



Origin of refugees

DRC **BURUNDI**

Site opened
Jan 1996

Partners



Key Indicators

CRUDE MORTALITY RATE

0.2 Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.4 Deaths/1,000 refugees/month

MATERNAL MORTALITY

1 Deaths

SKILLED BIRTH ATTENDANCE

95% of all births

GAM

2% Global Acute Malnutrition

SAM

0 Severe Acute Malnutrition


WATER SUPPLY

19 Liters/ person/ day



LATRINE COVERAGE

8 Refugees/communal toilets or latrines


COMMUNICABLE DISEASES




INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING			MORBIDITY		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Number of medical doctors	5	1 : < 50,000	✓	71	
Number of qualified nurses	101	1 : < 10,000	✓	4	
Number of community health workers	200	1 : < 1,000	✓	51	
ACCESS AND UTILIZATION			VACCINATION		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Consultations per trained clinician per day	40	< 50	✓	36%	< 95%
Health utilisation rate (new visits / refugee / year)	1.2	1 - 4	✓	95%	< 95%
Proportion of host population consultations	3%		✓	HIS	
MORTALITY			IN-PATIENT DEPARTMENT (IPD)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.2	< 0.75	✓	Average length of stay (days)	3.4
Under-five Mortality Rate (U5MR) (/1000/month)	0.4	< 1.5	✓	Case fatality rate	0.0
Infant Mortality Rate (IMR) (/1000 livebirths)	29.4	< 30	✓	Hospitalisation rate	50.9
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	7	< 20	✓		50 - 150
OUTBREAK ALERT AND RESPONSE					
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours	900%	100%	✓		
Number of outbreaks reported	2				

Reproductive Health

ANTENATAL CARE			FAMILY PLANNING		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Antenatal care coverage	73%	> 90%	✗	66%	≥ 30%
Coverage of antenatal tetanus vaccination	79%	> 95%	⚠	HIS	
DELIVERY CARE			SEXUAL AND GENDER-BASED VIOLENCE (SGBV)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of births attended by skilled personnel	95%	≥ 90%	✓	Total number of reported rapes	88
Proportion of births conducted by caesarean section	7%	5 - 15%	✓	Proportion of eligible rape survivors provided with PEP within 72 hours	87%
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	9%	< 15%	✓	Proportion of eligible rape survivors provided with ECP within 120 hours	55%
Proportion of teenage pregnancies	4%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	
Still birth rate (/ 1000 total births / month)	1			Number of genital ulcer diseases	41
Number of maternal deaths	1			Number of Pelvic Inflammatory diseases	17
Proportion of maternal deaths investigated within 48 hours	100%	100%	✓		

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD	PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV	83%	100%	✔ Number of PoCs receiving ART	748	
Condom distribution rate	0.12	> 0.5	✘ Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%	PMTCT coverage	90%	100% ⚠

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD	INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD
Prevalence of global acute malnutrition (6-59 months)	2%	< 20%	✔ Proportion of children 0-23 months who were put to the breast within one hour of birth	77%	> 80% ⚠
Prevalence of stunting (6-59 months)	43%	< 20%	✔ Proportion of infants aged between 0-5 months who received only breast milk during the previous day	75%	> 70% ✔
Prevalence of severe acute malnutrition (6 - 59 months)	0%	< 3%	✔ IYCF programmes targeting pregnant and lactating women established or maintained?		Yes
Prevalence of anaemia in children (6-59 months)	46%	< 20%	✘ MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD
Prevalence of anaemia in women of reproductive age (15-49 yrs)	25%	< 20%	✘ Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	85%	> 70% ✔

CMAM: SAM TREATMENT	INDICATOR	STANDARD	MAM TREATMENT	INDICATOR	STANDARD
Number of new severe acute admissions (6-59 months)	356		Number of new moderate acute admissions (6-59 months)	831	
Recovery rate	94%	> 75%	✔ Recovery rate	88%	> 75% ✔
Default rate	4%	< 0%	✘ Default rate	0%	< 15% ✔
Death rate	2%	< 10%	✔ Death rate	0%	< 3% ✔
Referral rate	1%		Referral rate	6%	

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD	WATER STAFFING	INDICATOR	STANDARD
Average number of litres of potable water available per person per day	19	< 20	⚠ Number of hygiene promoters	106	1 : < 20,000 ✔
Proportion of households collecting >= 15 litres per person per day	100%	> 95%	✔ SANITATION	INDICATOR	STANDARD
Number of persons of concern per water tap	89	< 80	⚠ Refugees per latrine/toilet	8	< 20 ✔
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5	100%	> 95%	✔ Proportion of households with drop-hole latrine or drop-hole toilet	44%	
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	0%	> 95%	✔ Proportion of households report defecating in a toilet	99%	> 85% ✔
Proportion of households collecting drinking water from protected water sources only	98%	> 95%	✘ Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	83%	> 90% ⚠
Proportion of households with sufficient daily water storage capacity	55%	> 80%			