



# Site report 2018 Kyangwali UGANDA



## Population

**92,308** refugees



### Origin of refugees

DRC SUDAN RWANDA  
BURUNDI

Site opened  
Jan 1968

## Partners



## Key Indicators

### CRUDE MORTALITY RATE

**0.3** Deaths/1,000 refugees/month

### U5 MORTALITY RATE

**0.7** Deaths/1,000 refugees/month

### MATERNAL MORTALITY

**6** Deaths

### SKILLED BIRTH ATTENDANCE

**93%** of all births

### GAM

**NC** Global Acute Malnutrition

### SAM

**NC** Severe Acute Malnutrition


### WATER SUPPLY

**14** Liters/ person/ day



### LATRINE COVERAGE

**5** Refugees/communal toilets or latrines


## COMMUNICABLE DISEASES




## INJURIES



## NON-COMMUNICABLE DISEASES



## MENTAL HEALTH





## HIV/AIDS

### PREVENTION & TREATMENT

Proportion of blood units screened for HIV

INDICATOR STANDARD

100%

Condom distribution rate

0.22 > 0.5

Proportion of TB patients tested for HIV

100%

### PREVENTION & TREATMENT

Number of PoCs receiving ART

INDICATOR STANDARD

808

✘ Proportion of HIV positive mothers who swallowed ARV during delivery

100%

PMTCT coverage

78% 100%



## Nutrition

### NUTRITION GENERAL

Prevalence of global acute malnutrition (6-59 months)

INDICATOR STANDARD

Not collected

Prevalence of stunting (6-59 months)

Not collected

Prevalence of severe acute malnutrition (6 - 59 months)

Not collected

Prevalence of anaemia in children (6-59 months)

Not collected

Prevalence of anaemia in women of reproductive age (15-49 yrs)

Not collected

### INFANT AND YOUNG CHILD FEEDING (IYCF)

Proportion of children 0-23 months who were put to the breast within one hour of birth

INDICATOR STANDARD

Not collected

Proportion of infants aged between 0-5 months who received only breast milk during the previous day

Not collected

IYCF programmes targeting pregnant and lactating women established or maintained?

Yes

### MICRONUTRIENT DEFICIENCY

Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)

INDICATOR STANDARD

77% > 70%



### CMAM: SAM TREATMENT

Number of new severe acute admissions (6-59 months)

INDICATOR STANDARD

1450

Recovery rate

83% > 75%



Default rate

11% < 0%



Death rate

1% < 10%



Referral rate

2%

### MAM TREATMENT

Number of new moderate acute admissions (6-59 months)

INDICATOR STANDARD

2690

Recovery rate

83% > 75%



Default rate

14% < 15%



Death rate

0% < 3%



Referral rate

2%

## Water, Sanitation and Hygiene (WASH)

### WATER

Average number of litres of potable water available per person per day

INDICATOR STANDARD

14 < 20



Proportion of households collecting >= 15 litres per person per day

39% > 95%



Number of persons of concern per water tap

218 < 80



Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual ( >= 0.1 mg/litre) and NTU <5

99% > 95%



Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml

64% > 95%



Proportion of households collecting drinking water from protected water sources only

92% > 95%



Proportion of households with sufficient daily water storage capacity

62% > 80%



### WATER STAFFING

Number of hygiene promoters

INDICATOR STANDARD

60 1 : < 20,000



### SANITATION

Refugees per latrine/toilet

INDICATOR STANDARD

5 < 20



Proportion of households with drop-hole latrine or drop-hole toilet

63%

Proportion of households report defecating in a toilet

61% > 85%



Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)

100% > 90%

