



Site report 2018 Rwamwanja UGANDA



Population

64,811 refugees



Origin of refugees

DRC

Site opened
Apr 2012

Partners



Key Indicators

CRUDE MORTALITY RATE

0.1 Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.2 Deaths/1,000 refugees/month

MATERNAL MORTALITY

2 Deaths

SKILLED BIRTH ATTENDANCE

95% of all births

GAM

NC Global Acute Malnutrition

SAM

NC Severe Acute Malnutrition


WATER SUPPLY

15 Liters/ person/ day



LATRINE COVERAGE

6 Refugees/communal toilets or latrines


COMMUNICABLE DISEASES




INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



HIV/AIDS

PREVENTION & TREATMENT

Proportion of blood units screened for HIV

INDICATOR STANDARD

100%

Condom distribution rate

0.49 > 0.5

Proportion of TB patients tested for HIV

100%

PREVENTION & TREATMENT

Number of PoCs receiving ART

INDICATOR STANDARD

422

⚠️ Proportion of HIV positive mothers who swallowed ARV during delivery

100%

PMTCT coverage

100% 100%



Nutrition

NUTRITION GENERAL

Prevalence of global acute malnutrition (6-59 months)

INDICATOR STANDARD

Not collected

Prevalence of stunting (6-59 months)

Not collected

Prevalence of severe acute malnutrition (6 - 59 months)

Not collected

Prevalence of anaemia in children (6-59 months)

Not collected

Prevalence of anaemia in women of reproductive age (15-49 yrs)

Not collected

INFANT AND YOUNG CHILD FEEDING (IYCF)

Proportion of children 0-23 months who were put to the breast within one hour of birth

INDICATOR STANDARD

Not collected

Proportion of infants aged between 0-5 months who received only breast milk during the previous day

Not collected

IYCF programmes targeting pregnant and lactating women established or maintained?

Yes

MICRONUTRIENT DEFICIENCY

Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)

INDICATOR STANDARD

80% > 70%



CMAM: SAM TREATMENT

Number of new severe acute admissions (6-59 months)

INDICATOR STANDARD

210

Recovery rate

84% > 75%



Default rate

14% < 0%



Death rate

0% < 10%



Referral rate

2%

MAM TREATMENT

Number of new moderate acute admissions (6-59 months)

INDICATOR STANDARD

434

Recovery rate

83% > 75%



Default rate

8% < 15%



Death rate

0% < 3%



Referral rate

5%

Water, Sanitation and Hygiene (WASH)

WATER

Average number of litres of potable water available per person per day

INDICATOR STANDARD

15 < 20



Proportion of households collecting >= 15 litres per person per day

64% > 95%



Number of persons of concern per water tap

298 < 80



Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5

100% > 95%



Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/100 ml

87% > 95%



Proportion of households collecting drinking water from protected water sources only

97% > 95%



Proportion of households with sufficient daily water storage capacity

30% > 80%



WATER STAFFING

Number of hygiene promoters

INDICATOR STANDARD

190 1 : < 20,000



SANITATION

Refugees per latrine/toilet

INDICATOR STANDARD

6 < 20



Proportion of households with drop-hole latrine or drop-hole toilet

83%

Proportion of households report defecating in a toilet

90% > 85%



Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)

0% > 90%

