



Site report 2018 Palorinya UGANDA



Population

119,071 refugees



Origin of refugees
SOUTH SUDAN

Site opened
Jan 1994

Partners



Key Indicators

CRUDE MORTALITY RATE

0.1 Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.1 Deaths/1,000 refugees/month

MATERNAL MORTALITY

2 Deaths

SKILLED BIRTH ATTENDANCE

97% of all births

GAM

NC Global Acute Malnutrition

SAM

NC Severe Acute Malnutrition


WATER SUPPLY

20 Liters/ person/ day



LATRINE COVERAGE

8 Refugees/communal toilets or latrines


COMMUNICABLE DISEASES




INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

| HEALTH STAFFING | | | MORBIDITY | | | |
|---|-----------|--------------|-----------------------------|--|-----------------|--|
| | INDICATOR | STANDARD | | INDICATOR | STANDARD | |
| Number of medical doctors | 6 | 1 : < 50,000 | ✓ | Incidence of malaria among children under 5 | 52 | |
| Number of qualified nurses | 65 | 1 : < 10,000 | ✓ | Incidence of watery diarrhoea among children under 5 | 13 | |
| Number of community health workers | 176 | 1 : < 1,000 | ✓ | Incidence of pneumonia among children under 5 | 25 | |
| ACCESS AND UTILIZATION | | | VACCINATION | | | |
| | INDICATOR | STANDARD | | INDICATOR | STANDARD | |
| Consultations per trained clinician per day | 49 | < 50 | ✓ | Incidence of pneumonia among over 5 | 11 | |
| Health utilisation rate (new visits / refugee / year) | 1.2 | 1 - 4 | ✓ | Tuberculosis success rate | 92% < 90% ✓ | |
| Proportion of host population consultations | 15% | | | Were any MDR/X-TB cases diagnosed among PoCs? | Yes No ✗ | |
| MORTALITY | | | IN-PATIENT DEPARTMENT (IPD) | | | |
| | INDICATOR | STANDARD | | INDICATOR | STANDARD | |
| Crude Mortality Rate (CMR) (/1000/month) | 0.1 | < 0.75 | ✓ | Full vaccination coverage | 123% < 95% ✓ | |
| Under-five Mortality Rate (U5MR) (/1000/month) | 0.1 | < 1.5 | ✓ | Measles vaccination coverage | Not collected | |
| Infant Mortality Rate (IMR) (/1000 livebirths) | 5.1 | < 30 | ✓ | Source of vaccination data | HIS | |
| Neonatal Mortality Rate (NNMR) (/1000 livebirths) | 3 | < 20 | ✓ | IN-PATIENT DEPARTMENT (IPD) | | |
| OUTBREAK ALERT AND RESPONSE | | | | INDICATOR | STANDARD | |
| | INDICATOR | STANDARD | | | | |
| Proportion of outbreaks investigated within 48 hours | | 100% | | Average length of stay (days) | 2.0 | |
| Number of outbreaks reported | 0 | | | Case fatality rate | 0.0 | |
| | | | | Hospitalisation rate | 48.2 50 - 150 ⚠ | |

Reproductive Health

| ANTENATAL CARE | | | FAMILY PLANNING | | | |
|--|-----------|----------|---|--|-------------|--|
| | INDICATOR | STANDARD | | INDICATOR | STANDARD | |
| Antenatal care coverage | 89% | > 90% | ⚠ | Contraceptive prevalence rate | 25% ≥ 30% ⚠ | |
| Coverage of antenatal tetanus vaccination | 96% | > 95% | ✓ | Source of contraceptive prevalence data | HIS | |
| DELIVERY CARE | | | SEXUAL AND GENDER-BASED VIOLENCE (SGBV) | | | |
| | INDICATOR | STANDARD | | INDICATOR | STANDARD | |
| Proportion of births attended by skilled personnel | 97% | ≥ 90% | ✓ | Total number of reported rapes | 130 | |
| Proportion of births conducted by caesarean section | 7% | 5 - 15% | ✓ | Proportion of eligible rape survivors provided with PEP within 72 hours | 90% 100% ⚠ | |
| Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs) | 7% | < 15% | ✓ | Proportion of eligible rape survivors provided with ECP within 120 hours | 60% 100% ✗ | |
| Proportion of teenage pregnancies | 5% | | | SEXUALLY TRANSMITTED INFECTIONS (STIS) | | |
| Still birth rate (/ 1000 total births / month) | 1 | | | Number of genital ulcer diseases | 205 | |
| Number of maternal deaths | 2 | | | Number of Pelvic Inflammatory diseases | 767 | |
| Proportion of maternal deaths investigated within 48 hours | 100% | 100% | ✓ | | | |

HIV/AIDS

PREVENTION & TREATMENT

Proportion of blood units screened for HIV

Condom distribution rate

Proportion of TB patients tested for HIV

INDICATOR

0.21

STANDARD

100%

> 0.5

100%

PREVENTION & TREATMENT

Number of PoCs receiving ART

✘ Proportion of HIV positive mothers who swallowed ARV during delivery

PMTCT coverage

INDICATOR

253

100%

97%

100%



Nutrition

NUTRITION GENERAL

Prevalence of global acute malnutrition (6-59 months)

Prevalence of stunting (6-59 months)

Prevalence of severe acute malnutrition (6 - 59 months)

Prevalence of anaemia in children (6-59 months)

Prevalence of anaemia in women of reproductive age (15-49 yrs)

INDICATOR

Not collected

Not collected

Not collected

Not collected

Not collected

STANDARD

INFANT AND YOUNG CHILD FEEDING (IYCF)

Proportion of children 0-23 months who were put to the breast within one hour of birth

Proportion of infants aged between 0-5 months who received only breast milk during the previous day

IYCF programmes targeting pregnant and lactating women established or maintained?

INDICATOR

Not collected

Not collected

Yes

INDICATOR

94%

STANDARD

> 70%



CMAM: SAM TREATMENT

Number of new severe acute admissions (6-59 months)

Recovery rate

Default rate

Death rate

Referral rate

INDICATOR

263

78%

6%

0%

14%

STANDARD

> 75%

< 0%

< 10%

MAM TREATMENT

Number of new moderate acute admissions (6-59 months)

✓ Recovery rate

✘ Default rate

✓ Death rate

Referral rate

INDICATOR

1555

84%

8%

0%

7%

STANDARD

> 75%

< 15%

< 3%



Water, Sanitation and Hygiene (WASH)

WATER

Average number of litres of potable water available per person per day

Proportion of households collecting >= 15 litres per person per day

Number of persons of concern per water tap

Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5

Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml

Proportion of households collecting drinking water from protected water sources only

Proportion of households with sufficient daily water storage capacity

INDICATOR

20

70%

125

100%

92%

73%

60%

STANDARD

< 20

> 95%

< 80

> 95%

> 95%

> 95%

> 80%



WATER STAFFING

Number of hygiene promoters

SANITATION

Refugees per latrine/toilet

✘ Proportion of households with drop-hole latrine or drop-hole toilet

✓ Proportion of households report defecating in a toilet

⚠ Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)

INDICATOR

160

8

47%

85%

0%

STANDARD

1 : < 20,000

< 20

> 85%

> 85%

> 90%

