


Site report 2018 Kiryandongo UGANDA



Population

57,639 refugees



Origin of refugees

SUDAN KENYA DRC BURUNDI

Site opened
Jan 1995

Partners



TPO Uganda

Key Indicators

CRUDE MORTALITY RATE

0.0 Deaths/1,000 refugees/month

U5 MORTALITY RATE

0.0 Deaths/1,000 refugees/month

MATERNAL MORTALITY

0 Deaths

SKILLED BIRTH ATTENDANCE

94% of all births

GAM

NC Global Acute Malnutrition

SAM

NC Severe Acute Malnutrition


WATER SUPPLY

16 Liters/ person/ day



LATRINE COVERAGE

4 Refugees/communal toilets or latrines


COMMUNICABLE DISEASES




INJURIES



NON-COMMUNICABLE DISEASES



MENTAL HEALTH



Public Health

HEALTH STAFFING			MORBIDITY		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Number of medical doctors	2	1 : < 50,000	✓	13	
Number of qualified nurses	33	1 : < 10,000	✓	11	
Number of community health workers	198	1 : < 1,000	✓	3	
ACCESS AND UTILIZATION			VACCINATION		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Consultations per trained clinician per day	34	< 50	✓	Incidence of pneumonia among over 5	1
Health utilisation rate (new visits / refugee / year)	0.8	1 - 4	⚠	Tuberculosis success rate	75% < 90% ⚠
Proportion of host population consultations	43%			Were any MDR/X-TB cases diagnosed among PoCs?	No No ✗
MORTALITY			IN-PATIENT DEPARTMENT (IPD)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Crude Mortality Rate (CMR) (/1000/month)	0.0	< 0.75	✓	Full vaccination coverage	52% < 95% ✗
Under-five Mortality Rate (U5MR) (/1000/month)	0.0	< 1.5	✓	Measles vaccination coverage	Not collected
Infant Mortality Rate (IMR) (/1000 livebirths)	1.1	< 30	✓	Source of vaccination data	HIS
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	1	< 20	✓	Average length of stay (days)	3.7
OUTBREAK ALERT AND RESPONSE			CASE FATALITY RATE		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0
Number of outbreaks reported	0			Hospitalisation rate	108.1 50 - 150 ✓

Reproductive Health

ANTENATAL CARE			FAMILY PLANNING		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Antenatal care coverage	68%	> 90%	✗	Contraceptive prevalence rate	1% ≥ 30% ✗
Coverage of antenatal tetanus vaccination	87%	> 95%	⚠	Source of contraceptive prevalence data	HIS
DELIVERY CARE			SEXUAL AND GENDER-BASED VIOLENCE (SGBV)		
	INDICATOR	STANDARD		INDICATOR	STANDARD
Proportion of births attended by skilled personnel	94%	≥ 90%	✓	Total number of reported rapes	11
Proportion of births conducted by caesarean section	3%	5 - 15%	✗	Proportion of eligible rape survivors provided with PEP within 72 hours	75% 100% ✗
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	3%	< 15%	✓	Proportion of eligible rape survivors provided with ECP within 120 hours	36% 100% ✗
Proportion of teenage pregnancies	11%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	
Still birth rate (/ 1000 total births / month)	0			Number of genital ulcer diseases	85
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	689
Proportion of maternal deaths investigated within 48 hours		100%			

HIV/AIDS

PREVENTION & TREATMENT

Proportion of blood units screened for HIV

INDICATOR STANDARD

100%

Condom distribution rate

0.68 > 0.5

Proportion of TB patients tested for HIV

100%

PREVENTION & TREATMENT

Number of PoCs receiving ART

INDICATOR STANDARD

389

✓ Proportion of HIV positive mothers who swallowed ARV during delivery

100%

PMTCT coverage

94% 100%



Nutrition

NUTRITION GENERAL

Prevalence of global acute malnutrition (6-59 months)

INDICATOR STANDARD

Not collected

Prevalence of stunting (6-59 months)

Not collected

Prevalence of severe acute malnutrition (6 - 59 months)

Not collected

Prevalence of anaemia in children (6-59 months)

Not collected

Prevalence of anaemia in women of reproductive age (15-49 yrs)

Not collected

INFANT AND YOUNG CHILD FEEDING (IYCF)

Proportion of children 0-23 months who were put to the breast within one hour of birth

INDICATOR STANDARD

Not collected

Proportion of infants aged between 0-5 months who received only breast milk during the previous day

Not collected

IYCF programmes targeting pregnant and lactating women established or maintained?

Yes

MICRONUTRIENT DEFICIENCY

Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)

INDICATOR STANDARD

71% > 70%



CMAM: SAM TREATMENT

Number of new severe acute admissions (6-59 months)

INDICATOR STANDARD

30

Recovery rate

94% > 75%

✓ Recovery rate

Default rate

6% < 0%

✗ Default rate

Death rate

0% < 10%

✓ Death rate

Referral rate

0%

MAM TREATMENT

Number of new moderate acute admissions (6-59 months)

INDICATOR STANDARD

139

✓ Recovery rate

100% > 75%



✗ Default rate

0% < 15%



✓ Death rate

0% < 3%



Referral rate

0%

Water, Sanitation and Hygiene (WASH)

WATER

Average number of litres of potable water available per person per day

INDICATOR STANDARD

16 < 20

Proportion of households collecting >= 15 litres per person per day

70% > 95%



Number of persons of concern per water tap

329 < 80



Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5

98% > 95%



Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/100 ml

100% > 95%



Proportion of households collecting drinking water from protected water sources only

100% > 95%



Proportion of households with sufficient daily water storage capacity

58% > 80%



WATER STAFFING

Number of hygiene promoters

INDICATOR STANDARD

200 1 : < 20,000



SANITATION

Refugees per latrine/toilet

INDICATOR STANDARD

4 < 20



✗ Proportion of households with drop-hole latrine or drop-hole toilet

73%

✓ Proportion of households report defecating in a toilet

70% > 85%



✓ Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)

100% > 90%

