

Site report 2018 Imvepi UGANDA

Population

53,958 refugees





Refugee Agency

Origin of refugees

SOUTH SUDAN

Site opened Jan 1995

Partners





Key Indicators

CRUDE MORTALITY RATE

Deaths/1,000 refugees/

GAM

Global Acute

U5 MORTALITY RATE

Deaths/1,000 refugees/

SAM

NC

Severe Acute Malnutrition

MATERNAL MORTALITY

Deaths

WATER SUPPLY

17

Liters/ person/ day

SKILLED BIRTH **ATTENDANCE**


95%


of all births

LATRINE COVERAGE


Refugees/communal toilets or latrines


COMMUNICABLE DISEASES





NON-COMMUNICABLE DISEASES





Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARD		
Number of medical doctors	1	1:<50,000	8	Incidence of malaria among children under 5	138			
Number of qualified nurses	20	1:<10,000	•	Incidence of watery diarrhoea among	24			
Number of community health workers	95	1:<1,000	•	children under 5				
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	34			
Consultations per trained clinician per day	50	< 50	A	Incidence of pneumonia among over 5	18			
Health utilisation rate (new visits / refugee /	2.1	1 - 4	•	Tuberculosis success rate	91%	< 90%	•	
year) Proportion of host population consultations	17%			Were any MDR/X-TB cases diagnosed among PoCs?	Yes	No	©	
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATO	OR STANI	DARD	
Crude Mortality Rate (CMR) (/1000/month)	0.1	< 0.75	②	Full vaccination coverage	81%	< 9	5%	×
Under-five Mortality Rate (U5MR) (/1000/month)	0.2	< 1.5	②	Measles vaccination coverage		Not collected	t	
Infant Mortality Rate (IMR) (/1000 livebirths)	32.5	< 30	Δ	Source of vaccination data	HIS			
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	29	< 20	8	IN-PATIENT DEPARTMENT (IPD)	INDICATO	STANI	DARD	
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	2.7			
Proportion of outbreaks investigated within 48 hours		100%		Case fatality rate	0.0			
Number of outbreaks reported	0			Hospitalisation rate	48.7	50 -	150	4

Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	96%	> 90%	②	Contraceptive prevalence rate	28%	≥ 30%	<u> </u>
Coverage of antenatal tetanus vaccination	98%	> 95%	②	Source of contraceptive prevalence data	HIS		
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	95%	≥ 90%	②	Total number of reported rapes	42		
Proportion of births conducted by caesarean section	13%	5 - 15%	②	Proportion of eligible rape survivors provided with PEP within 72 hours	79%	100%	8
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	7%	< 15%	②	Proportion of eligible rape survivors provided with ECP within 120 hours	75%	100%	8
Proportion of teenage pregnancies	4%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate (/ 1000 total births / month)	1			Number of genital ulcer diseases	56		
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	342		
Proportion of maternal deaths investigated within 48 hours		100%					

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD		PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%		Number of PoCs receving ART	570	
Condom distribution rate	0.64	> 0.5	②	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%		PMTCT coverage	100%	100%

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD		INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD	
Prevalence of global acute malnutrition (6-59 months)	Not	collected		Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected		
Prevalence of stunting (6-59 months)	Not	collected		Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected		
Prevalence of severe acute malnutrition (6 - 59 months)	Not	Not collected Not collected Not collected Not collected Not collected Not collected STANDARD 85 89% > 75%		IYCF programmes targeting pregnant and lactating women established or maintained?		Yes	
Prevalence of anaemia in children (6-59 months)	Not	collected		MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not	Not collected		Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	96%	> 70%	Ø
CMAM: SAM TREATMENT	INDICATOR	STANDARD		MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)	85			Number of new moderate acute admissions (6-59 months)	749		
Recovery rate	89%	> 75%	②	Recovery rate	81%	> 75%	②
Default rate	8%	< 0%	8	Default rate	10%	< 15%	②
Death rate	1%	< 10%	②	Death rate	0%	< 3%	②
Referral rate	0%			Referral rate	4%		

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	17	< 20	8	Number of hygiene promoters	152	1:<20,000	•
Proportion of households collecting >= 15 litres per person per day	75%	> 95%	A	SANITATION Refugees per latrine/toilet	INDICATOR 51	STANDARD < 20	8
Number of persons of concern per water tap	245	< 80	8	Proportion of households with drop-hole	55%	120	
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU	84%	> 95%	A	latrine or drop-hole toilet Proportion of households report defecating in a toilet	54%	> 85%	8
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	75%	> 95%	A	Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	75%	> 90%	A
Proportion of households collecting drinking water from protected water sources only	82%	> 95%	A				
Proportion of households with sufficient daily water storage capacity	94%	> 80%	•				