

Site report 2018 Nakivale UGANDA

Population

103,721 refugees







Origin of refugees

DRC SOMALIA RWANDA BURUNDI

Site opened Jan 1959

Partners







Key Indicators

CRUDE MORTALITY RATE

Deaths/1,000 refugees/

month

GAM

Global Acute Malnutrition U5 MORTALITY RATE

0.2

Deaths/1,000 refugees/

SAM

NC

Severe Acute Malnutrition MATERNAL MORTALITY

N

Deaths

WATER SUPPLY

17

Liters/ person/ day

SKILLED BIRTH ATTENDANCE


94% of all births


LATRINE COVERAGE

7


Refugees/communal toilets or latrines

COMMUNICABLE DISEASES





NON-COMMUNICABLE DISEASES



Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARD		
Number of medical doctors	3	1:<50,000	A	Incidence of malaria among children under 5	6			
Number of qualified nurses	75	1:<10,000	②	Incidence of watery diarrhoea among	24			
Number of community health workers	224	1:<1,000	②	children under 5				
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	29			
Consultations per trained clinician per day	44	< 50	②	Incidence of pneumonia among over 5	10			
Health utilisation rate (new visits / refugee /	1.0	1 - 4	Ø	Tuberculosis success rate	76%	< 90%	A	
year) Proportion of host population consultations	13%			Were any MDR/X-TB cases diagnosed among PoCs?	No	No	8	
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICAT	OR STAND	ARD	
Crude Mortality Rate (CMR) (/1000/month)	0.1	< 0.75	②	Full vaccination coverage	NR			
Under-five Mortality Rate (U5MR) (/1000/month)	0.2	< 1.5	②	Measles vaccination coverage		Not collected		
Infant Mortality Rate (IMR) (/1000 livebirths)	7.7	< 30	⊘	Source of vaccination data	HIS			
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	4	< 20	②	IN-PATIENT DEPARTMENT (IPD)	INDICAT	OR STAND	ARD	
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	2.1			
Proportion of outbreaks investigated within 48 hours	100%	100%	②	Case fatality rate	0.0			
Number of outbreaks reported	5			Hospitalisation rate	69.0	50 - 1	150	②

Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	58%	> 90%	8	Contraceptive prevalence rate	32%	≥ 30%	•
Coverage of antenatal tetanus vaccination	95%	> 95%	②	Source of contraceptive prevalence data	HIS		
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	94%	≥ 90%	⊘	Total number of reported rapes	56		
Proportion of births conducted by caesarean section	3%	5 - 15%	8	Proportion of eligible rape survivors provided with PEP within 72 hours	72%	100%	8
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	4%	< 15%	②	Proportion of eligible rape survivors provided with ECP within 120 hours	33%	100%	8
Proportion of teenage pregnancies	2%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate (/ 1000 total births / month)	1			Number of genital ulcer diseases	154		
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	495		
Proportion of maternal deaths investigated within 48 hours		100%					

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD		PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV	100%	100%	②	Number of PoCs receving ART	1260	
Condom distribution rate	0.19	> 0.5	8	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%		PMTCT coverage	95%	100%

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD		INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD	
Prevalence of global acute malnutrition (6-59 months)	Not collected			Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected		
Prevalence of stunting (6-59 months)	Not collected			Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected		
Prevalence of severe acute malnutrition (6 - 59 months)	Not collected			IYCF programmes targeting pregnant and lactating women established or maintained?	Yes		
Prevalence of anaemia in children (6-59 months)	Not collected			MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected			Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	97%	> 70%	Ø
CMAM: SAM TREATMENT	INDICATOR	STANDARD		MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)	163			Number of new moderate acute admissions (6-59 months)	600		
Recovery rate	92%	> 75%	②	Recovery rate	89%	> 75%	②
Default rate	6%	< 0%	8	Default rate	7%	< 15%	②
Death rate	1%	< 10%	②	Death rate	0%	< 3%	②
Referral rate	1%			Referral rate	0%		

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	17	< 20	8	Number of hygiene promoters	260	1:<20,000	②
Proportion of households collecting >= 15 litres per person per day	71%	> 95%	A	SANITATION Refugees per latrine/toilet	INDICATOR 7	STANDARD < 20	©
Number of persons of concern per water tap	1,020	< 80	8	Proportion of households with drop-hole latrine or drop-hole toilet	74%		
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5	92%	> 95%	A	Proportion of households report defecating in a toilet	96%	> 85%	•
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	70%	> 95%	A	Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	100%	> 90%	•
Proportion of households collecting drinking water from protected water sources only	87%	> 95%	A				
Proportion of households with sufficient daily water storage capacity	81%	> 80%	•				