

Site report 2018 Oruchinga UGANDA

Population

6,870 refugees







Origin of refugees

RVANDA

BURUNDI

Site opened Jan 1959

Partners



Key Indicators

CRUDE MORTALITY RATE

0.0

Deaths/1,000 refugees/ month

GAM

Global Acute Malnutrition **U5 MORTALITY RATE**

0.1

Deaths/1,000 refugees/

SAM

NC

Severe Acute Malnutrition MATERNAL MORTALITY

0

Deaths

WATER SUPPLY

23

Liters/ person/ day

SKILLED BIRTH ATTENDANCE

95%


of all births


LATRINE COVERAGE

9


Refugees/communal toilets or latrines


COMMUNICABLE DISEASES





NON-COMMUNICABLE DISEASES





Public Health

				MORPHRIT	INDICATOR	CTANDARD	
HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARD	
Number of medical doctors	1	1:<50,000	②	Incidence of malaria among children under 5	7		
Number of qualified nurses	12	1:<10,000	•	Incidence of watery diarrhoea among children under 5	19		
Number of community health workers	98	1:<1,000	②	Cilidren under 5			
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	24		
Consultations per trained clinician per day	25	< 50	•	Incidence of pneumonia among over 5	9		
Health utilisation rate (new visits / refugee /	1.1	1 - 4	•	Tuberculosis success rate	95%	< 90%	•
year)				Were any MDR/X-TB cases diagnosed among	No	No	8
Proportion of host population consultations	73%			PoCs?			
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATO	STANDAR	D
Crude Mortality Rate (CMR) (/1000/month)	0.0	< 0.75	②	Full vaccination coverage	72%	< 95%	8
Under-five Mortality Rate (U5MR) (/1000/month)	0.1	< 1.5	⊘	Measles vaccination coverage		Not collected	
Infant Mortality Rate (IMR) (/1000 livebirths)	5.0	< 30	②	Source of vaccination data	HIS		
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	0	< 20	②	IN-PATIENT DEPARTMENT (IPD)	INDICATO	STANDAR	D
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)	7.5		
Proportion of outbreaks investigated within 48 hours	100%	100%	②	Case fatality rate	0.0		
Number of outbreaks reported	1			Hospitalisation rate	62.2	50 - 150	②

Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	85%	> 90%	A	Contraceptive prevalence rate	38%	≥ 30%	•
Coverage of antenatal tetanus vaccination	99%	> 95%	②	Source of contraceptive prevalence data	HIS		
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	95%	≥ 90%	②	Total number of reported rapes	10		
Proportion of births conducted by caesarean section	1%	5 - 15%	8	Proportion of eligible rape survivors provided with PEP within 72 hours	70%	100%	8
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	2%	< 15%	②	Proportion of eligible rape survivors provided with ECP within 120 hours	67%	100%	8
Proportion of teenage pregnancies	2%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate (/ 1000 total births / month)	1			Number of genital ulcer diseases	22		
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	55		
Proportion of maternal deaths investigated within 48 hours		100%					

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD		PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%		Number of PoCs receving ART	184	
Condom distribution rate	0.88	> 0.5	②	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%		PMTCT coverage	98%	100%

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD		INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD	
Prevalence of global acute malnutrition (6-59 months)	Not collected			Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected		
Prevalence of stunting (6-59 months)	Not collected			Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected		
Prevalence of severe acute malnutrition (6 - 59 months)	Not collected			IYCF programmes targeting pregnant and lactating women established or maintained?	Yes		
Prevalence of anaemia in children (6-59 months)	Not collected			MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected			Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	95%	> 70%	Ø
CMAM: SAM TREATMENT	INDICATOR	STANDARD		MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)	32			Number of new moderate acute admissions (6-59 months)	132		
Recovery rate	94%	> 75%	②	Recovery rate	94%	> 75%	②
Default rate	6%	< 0%	8	Default rate	4%	< 15%	②
Death rate	0%	< 10%	②	Death rate	0%	< 3%	②
Referral rate	0%			Referral rate	2%		

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	23	< 20	•	Number of hygiene promoters	40	1:<20,000	•
Proportion of households collecting >= 15 litres per person per day		> 95%		SANITATION Refugees per latrine/toilet	INDICATOR 9	STANDARD < 20	•
Number of persons of concern per water tap	258	< 80	8	Proportion of households with drop-hole latrine or drop-hole toilet	78%		
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5	76%	> 95%	A	Proportion of households report defecating in a toilet	97%	> 85%	•
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	100%	> 95%	⊘	Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	100%	> 90%	•
Proportion of households collecting drinking water from protected water sources only	92%	> 95%	A				
Proportion of households with sufficient daily water storage capacity	73%	> 80%	A				