
Site report 2018 Basateen YEMEN





Origin of refugees SOMALIA ETHIOPIA SYRIAN ERITREA

Site opened Jan 1991

Partners



Key Indicators

SKILLED BIRTH CRUDE MORTALITY RATE **U5 MORTALITY RATE** MATERNAL MORTALITY ATTENDANCE 0.1 Deaths/1,000 refugees/ Deaths/1,000 refugees/ 98% of all births 0.40 Deaths month month GAM SAM WATER SUPPLY LATRINE COVERAGE NC Global Acute NC Severe Acute 20 Refugees/communal toilets or latrines Liters/ person/ day Malnutrition Malnutrition COMMUNICABLE DISEASES **NON-COMMUNICABLE DISEASES** 37.5 % 30.5 % Cardiovascular disease URTI 18.1 % 10.9 % Watery diarrhoea Other 29.4 % 6.5 % 7.0 % URTI 17.4 % 32.0 % Gynaecological Watery diarrhoea disorders Total 44.3 % 13.1 % Consultations 6.8 % Other 8,333 **MENTAL HEALTH** INJURIES 4.1 % 30.5 % 42.3 % Severe emotior Other disor 30.4 % Psychotic disorder 4.1 % Other

8.5 % Epilepsy seizures

26.5 %

Medicallv

unexplained

somatic complaint

Public Health

HEALTH STAFFING	INDICATOR	STANDARD		MORBIDITY	INDICATOR	STANDARD	
Number of medical doctors	7	1 : < 50,000	0	Incidence of malaria among children under 5	0		
Number of qualified nurses	6	1 : < 10,000	0	Incidence of watery diarrhoea among	13		
Number of community health workers	16	1 : < 1,000	0	children under 5			
ACCESS AND UTILIZATION	INDICATOR	STANDARD		Incidence of pneumonia among children under 5	2		
Consultations per trained clinician per day	13	< 50	0	Incidence of pneumonia among over 5	0		
Health utilisation rate (new visits / refugee /	0.2	1 - 4	8	Tuberculosis success rate	90%	< 90%	O
year) Proportion of host population consultations	68%			Were any MDR/X-TB cases diagnosed among PoCs?	No	No	8
MORTALITY	INDICATOR	STANDARD		VACCINATION	INDICATOR	STANDAR	D
Crude Mortality Rate (CMR) (/1000/month)	0.1	< 0.75	0	Full vaccination coverage	8%	< 95%	8
Under-five Mortality Rate (U5MR) (/1000/month)	0.4	< 1.5	0	Measles vaccination coverage	13%	< 95%	8
Infant Mortality Rate (IMR) (/1000 livebirths)	49.2	< 30	8	Source of vaccination data	HIS		
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	18	< 20	0	IN-PATIENT DEPARTMENT (IPD)	INDICATOR	STANDAR	D
OUTBREAK ALERT AND RESPONSE	INDICATOR	STANDARD		Average length of stay (days)			
Proportion of outbreaks investigated within 48 hours	100%	100%	0	Case fatality rate			
Number of outbreaks reported	30			Hospitalisation rate	0.0	50 - 150	8

Reproductive Health

ANTENATAL CARE	INDICATOR	STANDARD		FAMILY PLANNING	INDICATOR	STANDARD	
Antenatal care coverage	0%	> 90%	⊗	Contraceptive prevalence rate	10%	≥ 30%	⊗
Coverage of antenatal tetanus vaccination	0%	> 95%	8	Source of contraceptive prevalence data	HIS		
DELIVERY CARE	INDICATOR	STANDARD		SEXUAL AND GENDER-BASED VIOLENCE (SGBV)	INDICATOR	STANDARD	
Proportion of births attended by skilled personnel	98%	≥ 90%	0	Total number of reported rapes	9		
Proportion of births conducted by caesarean section	0%	5 - 15%	8	Proportion of eligible rape survivors provided with PEP within 72 hours	200%	100%	⊘
Proportion of newborn infants with low birth weight (<2500 gs) (weighed within 72 hrs)	0%	< 15%	0	Proportion of eligible rape survivors provided with ECP within 120 hours	100%	100%	⊘
Proportion of teenage pregnancies	0%			SEXUALLY TRANSMITTED INFECTIONS (STIS)	INDICATOR	STANDARD	
Still birth rate (/ 1000 total births / month)	2			Number of genital ulcer diseases	9		
Number of maternal deaths	0			Number of Pelvic Inflammatory diseases	0		
Proportion of maternal deaths investigated within 48 hours		100%					

HIV/AIDS

PREVENTION & TREATMENT	INDICATOR	STANDARD	PREVENTION & TREATMENT	INDICATOR	STANDARD
Proportion of blood units screened for HIV		100%	Number of PoCs receving ART	20	
Condom distribution rate	0.01	> 0.5	Proportion of HIV positive mothers who swallowed ARV during delivery		100%
Proportion of TB patients tested for HIV		100%	PMTCT coverage	99%	100% 🛕

Nutrition

NUTRITION GENERAL	INDICATOR	STANDARD		INFANT AND YOUNG CHILD FEEDING (IYCF)	INDICATOR	STANDARD	
Prevalence of global acute malnutrition (6-59 months)	Not collected			Proportion of children 0-23 months who were put to the breast within one hour of birth	Not collected		
Prevalence of stunting (6-59 months)	Not collected			Proportion of infants aged between 0-5 months who received only breast milk during the previous day	Not collected		
Prevalence of severe acute malnutrition (6 - 59 months)	Not collected			IYCF programmes targeting pregnant and lactating women established or maintained?	Yes		
Prevalence of anaemia in children (6-59 months)	Not collected			MICRONUTRIENT DEFICIENCY	INDICATOR	STANDARD	
Prevalence of anaemia in women of reproductive age (15-49 yrs)	Not collected			Percentage coverage of special nutrition product for prevention of undernutrition in young children (LNS/MNP/FBF)	Not collected		
CMAM: SAM TREATMENT	INDICATOR	STANDARD		MAM TREATMENT	INDICATOR	STANDARD	
Number of new severe acute admissions (6-59 months)	79			Number of new moderate acute admissions (6-59 months)	103		
Recovery rate	100%	> 75%	0	Recovery rate	100%	> 75%	0
Default rate	0%	< 0%	8	Default rate	0%	< 15%	v
Death rate	0%	< 10%	0	Death rate	0%	< 3%	v
Referral rate	0%			Referral rate	0%		

Water, Sanitation and Hygiene (WASH)

WATER	INDICATOR	STANDARD		WATER STAFFING	INDICATOR	STANDARD	
Average number of litres of potable water available per person per day	20	< 20	0	Number of hygiene promoters	15	1:<20,000	0
Proportion of households collecting >= 15	Not collected			SANITATION	INDICATOR	STANDARD	
litres per person per day				Refugees per latrine/toilet	0	< 20	0
Number of persons of concern per water tap	Not collected			Proportion of households with drop-hole latrine or drop-hole toilet	Not collected		
Proportion of tests at chlorinated water collection points indicating a measurable chlorine residual (>= 0.1 mg/litre) and NTU <5	Not collected			Proportion of households report defecating in a toilet	Not collected		
Proportion of tests at unchlorinated water collection points showing 0 Faecal Coliforms/ 100 ml	Not collected			Proportion of communal latrines compliant with UNHCR standards (cleanable slabs, privacy & structural safety)	Not	collected	
Proportion of households collecting drinking water from protected water sources only	Not	collected					
Proportion of households with sufficient daily water storage capacity	Not	collected					